



DEPARTMENT OF MATHEMATICS

UNIT-V

LAPLACE TRANSFORMS

INTRODUCTION:

Laplace Transformation, named after a great French Mathematician Pierre Simon De Laplace (1749-1827) who used such transformations in the "Theory of probability".

Uses of Laplace Transformation:

1. It is used to find the solution of linear differential equations - ordinary as well as partial.
2. It helps in solving the differential equation with boundary values without finding the general solution and then finding the values of the arbitrary constants.

Transformation:

A transformation is an operation which converts a mathematical expression to a different but equivalent form.

Laplace Transformation: Definition:

Let $f(t)$ be a function of t defined for $t > 0$. Then the Laplace transform of $f(t)$, denoted by $\mathcal{L}\{f(t)\}$ or $F(s)$ is defined by,

$$\mathcal{L}[f(t)] = \int_0^{\infty} e^{-st} f(t) dt = F(s)$$

Provided the integral exists.



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Approved by AICTE, New Delhi, Affiliated to Anna University, Chennai

Accredited by NAAC-UGC with 'A++' Grade (Cycle III) & amp;

Accredited by NBA (B.E - CSE, EEE, ECE, Mech & amp; B.Tech.IT)

COIMBATORE-641 035, TAMIL NADU



DEPATMENT OF MATHEMATICS

Conditions for existence of Laplace transform:

- (i) $f(t)$ should be continuous or piecewise continuous in the given closed interval $[a, b]$ where $a > 0$.
- (ii) $f(t)$ should be of exponential order.

Exponential order:

A function $f(t)$ is said to be of exponential order if,

$$\lim_{t \rightarrow \infty} e^{-st} f(t) = 0.$$

Example:

1. t^2 is of exponential order.

$$\lim_{t \rightarrow \infty} e^{-st} f(t) = \lim_{t \rightarrow \infty} e^{-st} t^2$$

$$= \lim_{t \rightarrow \infty} \frac{t^2}{e^{st}} \quad \left[\frac{\infty}{\infty} \text{ Indeterminate form} \right]$$

$$= \lim_{t \rightarrow \infty} \frac{2t}{s e^{st}} \quad \text{Apply L'Hospital's rule}$$

$$= \lim_{t \rightarrow \infty} \frac{2}{s^2 e^{st}} = \frac{2}{\infty} = 0$$

2. $e t^2$ is not of exponential order.

$$\lim_{t \rightarrow \infty} e^{-st} f(t) = \lim_{t \rightarrow \infty} e^{-st} e t^2$$

$$= \lim_{t \rightarrow \infty} \frac{t^2}{e^{st-1}}$$

$$= e^{\infty} = \infty$$

$\therefore e t^2$ is not of exponential order.



DEPARTMENT OF MATHEMATICS

Transforms of elementary functions :

① $L(1) = \frac{1}{s}$ where $s > 0$

Proof :

$$L\{f(t)\} = \int_0^{\infty} e^{-st} f(t) dt$$

$$L(1) = \int_0^{\infty} e^{-st} \cdot 1 dt$$

$$= \left[\frac{e^{-st}}{-s} \right]_0^{\infty}$$

$$= -\frac{1}{s} (0 - 1) = \frac{1}{s}$$

$$\boxed{L(1) = \frac{1}{s}}$$

② $L(k) = \frac{k}{s}$

③ $L(t) = \frac{1!}{s^2}$

$$L(t) = \int_0^{\infty} e^{-st} \cdot t dt$$

$$= \left[\frac{te^{-st}}{-s} + \frac{e^{-st}}{s^2} \right]_0^{\infty}$$

$$\boxed{L(t) = \frac{1!}{s^2}}$$

④ $L(t^2) = \frac{2!}{s^3}$

⑤ $L(t^n) = \frac{n!}{s^{n+1}}$ if $s > 0$ & $n > -1$



DEPARTMENT OF MATHEMATICS

$$L(t^n) = \int_0^{\infty} e^{-st} t^n dt$$

$$\text{put } x = st \Rightarrow dx = s dt \quad \frac{1}{s} = (1) dt$$

$$\frac{dx}{s} = dt$$

$$L(t^n) = \int_0^{\infty} e^{-x} \left(\frac{x}{s}\right)^n \frac{dx}{s}$$

$$= \int_0^{\infty} e^{-x} \frac{x^n}{s^{n+1}} dx$$

$$= \frac{1}{s^{n+1}} \int_0^{\infty} e^{-x} x^n dx$$

$$L(t^n) = \frac{\Gamma_{n+1}}{s^{n+1}} = \frac{n!}{s^{n+1}}$$

$$(6) L(e^{at}) = \frac{1}{s-a} \text{ if } s-a > 0.$$

$$L(e^{at}) = \int_0^{\infty} e^{-st} e^{at} dt$$

$$= \int_0^{\infty} e^{-(s-a)t} dt$$

$$= \left[\frac{e^{-(s-a)t}}{-(s-a)} \right]_0^{\infty}$$

$$L(e^{at}) = \frac{1}{s-a} \text{ if } s-a > 0$$

$$(7) L(e^{-at}) = \frac{1}{s+a} \text{ if } s+a > 0$$

$$L(e^{-at}) = \int_0^{\infty} e^{-st} e^{-at} dt$$



DEPARTMENT OF MATHEMATICS

$$= \int_0^{\infty} e^{-(s+a)t} dt \quad (10 \text{ dec } 20) \text{ L. bnif } 01 \quad (1)$$

$$L(e^{-at}) = \frac{1}{s+a} \quad \text{if } s+a > 0$$

(8) To find $L(\cos at)$ & $L(\sin at)$:

We know $e^{i\theta} = \cos \theta + i \sin \theta$:

$$\begin{aligned} L(e^{iat}) &= \frac{1}{s-ia} \\ &= \frac{1}{s-ia} \cdot \frac{s+ia}{s+ia} = \frac{s+ia}{s^2+a^2} \\ &= \frac{s}{s^2+a^2} + i \frac{a}{s^2+a^2} \end{aligned}$$

$$L(\cos at + i \sin at) = \frac{s}{s^2+a^2} + i \frac{a}{s^2+a^2} \quad (1)$$

Equating real & imaginary parts,

$$\begin{aligned} L(\cos at) &= \frac{s}{s^2+a^2} \\ L(\sin at) &= \frac{a}{s^2+a^2} \end{aligned}$$

(9) To find $L(\sinh at)$:

$$L[\sinh at] = L\left(\frac{e^{at} - e^{-at}}{2}\right)$$

$$= \frac{1}{2} L(e^{at}) - \frac{1}{2} L(e^{-at})$$

$$= \frac{1}{2} \left[\frac{1}{s-a} - \frac{1}{s+a} \right]$$

$$= \frac{1}{2} \left(\frac{2a}{s^2-a^2} \right)$$

$$L(\sinh at) = \frac{a}{s^2-a^2} \quad \text{for } s^2 > a^2$$