



DEPARTMENT OF MATHEMATICS

UNIT- IV TESTING OF HYPOTHESIS

TEST OF SIGNIFICANCE OF SMALL SAMPLES!

STUDENT'S T-TEST :

TEST FOR SINGLE MEAN:

Null hypothesis : $H_0: \mu = \mu_0$

Test statistic, $t = \frac{\bar{x} - \mu}{s/\sqrt{n-1}}$ if SD is given.

$t = \frac{\bar{x} - \mu}{s/\sqrt{n}}$ if SD is not given.

To find s :

$$s^2 = \frac{\sum (x - \bar{x})^2}{n-1}$$

Degrees of freedom: $v = n-1$

NOTE: Confidence limit: $\bar{x} \pm t_{\alpha} \frac{s}{\sqrt{n-1}}$

- 1) A random sample of 10 boys had the following IQ's. 70, 120, 110, 101, 88, 88, 95, 98, 107, 100. Do these data support the assumption of a population mean IQ's of 100? Find a reasonable range to which most of the mean IQ's value of sample 10 boys.

Soln: given: $n = 10, \mu = 100$

$$\bar{x} = \frac{70 + 120 + 110 + 101 + 88 + 88 + 95 + 98 + 107 + 100}{10}$$

$$= 97.2$$



DEPARTMENT OF MATHEMATICS

UNIT- IV TESTING OF HYPOTHESIS

To find s :
$$s^2 = \frac{\sum (x - \bar{x})^2}{n-1}$$

$$x : 70 \quad 120 \quad 110 \quad 101 \quad 88 \quad 83 \quad 95 \quad 98 \quad 107 \quad 100$$

$$x - \bar{x} : -27.2 \quad 22.8 \quad 12.8 \quad 3.8 \quad -9.2 \quad -14.2 \quad -2.2 \quad 0.8 \quad 9.8 \quad 2.8$$

$$(x - \bar{x})^2 : 739.84 \quad 519.84 \quad 163.84 \quad 14.44 \quad 84.64 \quad 201.64 \quad 4.84 \quad 0.64 \quad 96.04 \quad 7.84$$

$$\sum (x - \bar{x})^2 = 1833.6$$

$$\therefore s^2 = \frac{\sum (x - \bar{x})^2}{n-1} = \frac{1833.6}{10-1}$$

$$= 203.73$$

$$\Rightarrow s = 14.27$$

Step 1: Formulating H_0 and H_1 :

$$H_0 : \mu = 100$$

$$H_1 : \mu \neq 100 \text{ (two tailed test)}$$

Step 2: Los. at $\alpha = 5\% = 0.05$

Step 3: Test statistic, $t = \frac{\bar{x} - \mu}{s/\sqrt{n}}$

$$= \frac{97.2 - 100}{14.27/\sqrt{10}}$$

$$= -0.62$$

$$|t| = 0.62$$



DEPARTMENT OF MATHEMATICS

UNIT- IV TESTING OF HYPOTHESIS

Step 4: t_{tab} for degree of freedom, $\nu = n - 1$
 $\nu = 10 - 1 = 9$

(a) $t_{tab} = 2.262 (t_{\alpha})$

Step 5: Conclusion: $t = 0.62 < 2.262 = t_{\alpha}$

$\therefore H_0$ is accepted at 5% LOS.

(a) The population mean IQ's is 100.

Confidence limit:

$$\mu = \bar{x} \pm t_{\alpha} \frac{s}{\sqrt{n-1}}$$

$$= 97.2 \pm 2.262 \times \frac{14.27}{\sqrt{10-1}}$$

$$= 97.2 \pm 10.759$$

$$= 107.95, 86.45$$

3) The weights of 10 peoples of a locality are found to be 70, 67, 62, 68, 61, 68, 70, 64, 64, 66 kg. It is reasonable to believe that the average weights of people locality greater than 64 kg. test at 5% LOS.

Soln: Given: $n = 10, \mu = 64$

$$\bar{x} = \frac{70 + 67 + 62 + 68 + 61 + 68 + 70 + 64 + 64 + 66}{10}$$

$$\bar{x} = 66$$



DEPARTMENT OF MATHEMATICS

UNIT- IV TESTING OF HYPOTHESIS

To find s :

$$s^2 = \frac{\sum (x - \bar{x})^2}{n-1}$$

x :	70	67	62	68	61	68	70	64	64	66
$x - \bar{x}$:	4	1	-4	2	-5	2	4	-2	-2	0
$(x - \bar{x})^2$:	16	1	16	4	25	4	16	4	4	0

$$\sum (x - \bar{x})^2 = 90$$

$$\therefore s^2 = \frac{\sum (x - \bar{x})^2}{n-1} = \frac{90}{10-1} = 10$$

$$s = 3.16$$

Step 1: Formulating H_0 and H_1 :

$$H_0: \mu = 64$$

$$H_1: \mu > 64 \text{ (one tailed test - right)}$$

Step 2: LOS at $\alpha = 5\%$.

Step 3: Test statistic, $t = \frac{\bar{x} - \mu}{s/\sqrt{n}}$

$$= \frac{66 - 64}{3.16/\sqrt{10}}$$

$$= 2.02$$

Step 4: t_{tab} for degree of freedom, $\nu = n-1$

$$= 10-1$$

$$= 9$$

As $t_{tab}: t_{\alpha} = 1.833$ (at two tailed at 10%.)



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore- 35



DEPARTMENT OF MATHEMATICS

UNIT- IV TESTING OF HYPOTHESIS

steps : Conclusion: $t = 2.02 > 1.833 = t_{\alpha}$
 $\therefore H_0$ is rejected at 5% LOS.
(a) the avg. weight of people locality is greater than 64 kg.



DEPARTMENT OF MATHEMATICS

UNIT- IV TESTING OF HYPOTHESIS

TEST FOR DIFFERENCE OF MEAN:

Null hypothesis; $H_0: \mu_1 = \mu_2$

$$\text{Test statistics, } t = \frac{\bar{x}_1 - \bar{x}_2}{s \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}$$

$$\text{where } s^2 = \frac{n_1 s_1^2 + n_2 s_2^2}{n_1 + n_2 - 2} \quad (\text{or}) \quad s^2 = \frac{\sum (x_1 - \bar{x}_1)^2 + \sum (x_2 - \bar{x}_2)^2}{n_1 + n_2 - 2}$$

Degree of freedom; $v = n_1 + n_2 - 2$.

1) In a test examination given to two groups of students, the marks obtained were as follows:

Group I : 18 20 36 50 49 36 34 49 41

Group II : 29 28 26 35 30 44 46

Examine whether the significance of difference between the average marks secured by the students of the above two groups.



DEPARTMENT OF MATHEMATICS

UNIT- IV TESTING OF HYPOTHESIS

Soln: Given: Group I: $n_1 = 9$

Group II: $n_2 = 7$

$$\text{Now } \bar{x}_1 = \frac{18 + 20 + 36 + 50 + 49 + 36 + 34 + 49 + 41}{9} = 34$$

$$\bar{x}_2 = \frac{29 + 28 + 26 + 35 + 30 + 44 + 46}{7} = 34$$

x_1	$x_1 - \bar{x}_1$	$(x_1 - \bar{x}_1)^2$	x_2	$(x_2 - \bar{x}_2)$	$(x_2 - \bar{x}_2)^2$
18	-19	361	29	-5	25
20	-14	289	28	-6	36
36	-1	1	26	-8	64
50	13	169	35	1	1
49	12	144	30	-4	16
36	-1	1	44	10	100
34	-3	9	46	12	144
49	12	144			
41	4	16			

$$\sum (x_1 - \bar{x}_1)^2 = 1134$$

$$\sum (x_2 - \bar{x}_2)^2 = 386$$

$$\text{Now } S^2 = \frac{\sum (x_1 - \bar{x}_1)^2 + \sum (x_2 - \bar{x}_2)^2}{n_1 + n_2 - 2}$$

$$= \frac{1134 + 386}{9 + 7 - 2} = 108.57$$

$$\Rightarrow S = 10.42$$



DEPARTMENT OF MATHEMATICS

UNIT- IV TESTING OF HYPOTHESIS

Step 1: Formulating H_0 and H_1

$$H_0: \mu_1 = \mu_2$$

$$H_1: \mu_1 \neq \mu_2 \text{ (two tailed test)}$$

Step 2: LOS at $\alpha = 5\%$.

Step 3: Test statistic, $t = \frac{\bar{x}_1 - \bar{x}_2}{S \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}$

$$= \frac{37 - 34}{10.42 \sqrt{\frac{1}{9} + \frac{1}{7}}}$$
$$= 0.5413$$

Step 4: t_{tab} for degrees of freedom, $\nu = n_1 + n_2 - 2$

$$= 9 + 4 - 2$$

$$\nu = 14$$

$$(iv) t_{tab} = (t_{\alpha}) = 2.145$$

Step 5: Conclusion: $t = 0.5413 < 2.145 = t_{\alpha}$.

$\therefore H_0$ is accepted at 5% LOS.

\therefore there is no significant difference in the avg. marks of the two groups of students.



DEPARTMENT OF MATHEMATICS

UNIT- IV TESTING OF HYPOTHESIS

2) A samples of two types of electric bulbs were tested for length of life and the following data were obtained.

Samples	size	mean	SD
I	8	1134	35
II	7	1024	40

Test at 5%

Soln: Given:

sample I : $n_1 = 8$, $\bar{x}_1 = 1134$, $s_1 = 35$

sample II : $n_2 = 7$, $\bar{x}_2 = 1024$, $s_2 = 40$

step 1: Formulating H_0 and H_1 .

$$H_0: \mu_1 = \mu_2$$

$$H_1: \mu_1 \neq \mu_2 \text{ (two tailed test)}$$

step 2: LOS at $\alpha = 5\%$.

step 3: Test statistic, $t = \frac{\bar{x}_1 - \bar{x}_2}{S \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}$

$$\text{Now } S^2 = \frac{n_1 s_1^2 + n_2 s_2^2}{n_1 + n_2 - 2}$$

$$= \frac{8(35)^2 + 7(40)^2}{8 + 7 - 2}$$

$$= \frac{8(1225) + 7(1600)}{13}$$

$$= 1615.38$$

$$S = 40.19$$



DEPARTMENT OF MATHEMATICS

UNIT- IV TESTING OF HYPOTHESIS

$$t = \frac{1134 - 1024}{40.19 \sqrt{\frac{1}{8} + \frac{1}{7}}}$$
$$= \frac{110}{20.8} = 5.288$$

step 4: t_{tab} for degrees of freedom, $\nu = n_1 + n_2 - 2$

$$= 8 + 7 - 2$$
$$= 13$$

(ii) $t_{tab}: t_{\alpha} = 2.160$

step 5: Conclusion: $t = 5.288 > 2.160 = t_{\alpha}$

$\therefore H_0$ is rejected at 5%.