



## SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Approved by AICTE, New Delhi, Affiliated to Anna University, Chennai

Accredited by NAAC-UGC with 'A++' Grade (Cycle III) &

Accredited by NBA (B.E - CSE, EEE, ECE, Mech & B.Tech.IT)

COIMBATORE-641 035, TAMIL NADU



### ODD AND EVEN FUNCTIONS:


#### EVEN FUNCTION:

Let  $f(x)$  be defined in  $(-1, 1)$   
If  $f(-x) = f(x)$ , then  $f(x)$  is an even function.

Note:

1. The graph of even function is symmetrical about y-axis
2.  $\int_{-1}^1 f(x) dx = 2 \int_0^1 f(x) dx$  if  $f(x)$  is even.
3. Sum of two even functions is also an even function.
4. Product of two even functions is also an even function.

#### Graph of an Even Function:





## SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Approved by AICTE, New Delhi, Affiliated to Anna University, Chennai

Accredited by NAAC-UGC with 'A++' Grade (Cycle III) &

Accredited by NBA (B.E - CSE, EEE, ECE, Mech & B.Tech.IT)

COIMBATORE-641 035, TAMIL NADU


### ODD FUNCTION:

Let  $f(x)$  be defined in  $(-a, a)$ .  
If  $f(-x) = -f(x)$ , then  $f(x)$  is an odd function.

#### Note:

1. The graph of odd function is symmetric about origin.
2.  $\int_{-a}^a f(x) dx = 0$  if  $f(x)$  is odd.
3. Sum of two odd function is also an odd function.
4. Product of two odd function is an even function.

### Graph of an Odd function:





## SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Approved by AICTE, New Delhi, Affiliated to Anna University, Chennai

Accredited by NAAC-UGC with 'A++' Grade (Cycle III) &

Accredited by NBA (B.E - CSE, EEE, ECE, Mech & B.Tech.IT)

COIMBATORE-641 035, TAMIL NADU

### PROBLEMS UNDER ODD AND EVEN FUNCTIONS:

1) Find the Fourier series for  $f(x) = x^2$  in  $(-1, 1)$  and hence deduce

$$(i) \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots = \frac{\pi^2}{6}$$

$$(ii) \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \dots = \frac{\pi^2}{12}$$

Solution:

$f(x)$  is defined in  $(-1, 1)$

In  $(-1, 1)$ , check whether  $f(x)$  is even / odd.

$$f(x) = x^2$$

$$f(-x) = (-x)^2 = x^2 = f(x)$$

$$f(x) = f(-x)$$

Hence,  $f(x)$  is an even function.

$$\therefore b_n = 0.$$

The Fourier series of  $f(x)$  in  $(-1, 1)$  is given by

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos \frac{n\pi x}{l} \quad \text{--- (1)}$$

$$a_0 = \frac{2}{l} \int_0^l f(x) dx$$

$$= \frac{2}{1} \int_0^1 x^2 dx$$

$$= \frac{2}{1} \left[ \frac{x^3}{3} \right]_0^1$$



# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Approved by AICTE, New Delhi, Affiliated to Anna University, Chennai

Accredited by NAAC-UGC with 'A++' Grade (Cycle III) &

Accredited by NBA (B.E - CSE, EEE, ECE, Mech & B.Tech.IT)

COIMBATORE-641 035, TAMIL NADU



$$= \frac{2}{1} \left( \frac{1^3}{3} \right)$$

$$a_0 = \frac{21^2}{3}$$

$$a_n = \frac{2}{1} \int_0^1 f(x) \cos \frac{n\pi x}{1} dx$$

$$= \frac{2}{1} \int_0^1 x^2 \cos \frac{n\pi x}{1} dx$$

$$= \frac{2}{1} \left[ (x^2) \left( \frac{\sin \frac{n\pi x}{1}}{\frac{n\pi}{1}} \right) - (2x) \left( \frac{-\cos \frac{n\pi x}{1}}{\left( \frac{n\pi}{1} \right)^2} \right) + (2) \left( \frac{-\sin \frac{n\pi x}{1}}{\left( \frac{n\pi}{1} \right)^3} \right) \right]_0^1$$

$$= \frac{4}{1} \left( \frac{1}{n\pi} \right)^2 \left[ x \cos \frac{n\pi x}{1} \right]_0^1$$

$$= \frac{41}{n^2\pi^2} [1 \cos n\pi - 0 \cos 0]$$

$$a_n = \frac{41^2 (-1)^n}{n^2\pi^2}$$

Substitute  $a_0, a_n$  in ①

$$f(x) = \frac{1^2}{3} + \sum_{n=1}^{\infty} \frac{41^2 (-1)^n}{n^2\pi^2} \cos \frac{n\pi x}{1} \quad \text{--- ②}$$

$$i) \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots = \frac{\pi^2}{6}$$

Put  $x=1$  in ②, we get





# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Approved by AICTE, New Delhi, Affiliated to Anna University, Chennai

Accredited by NAAC-UGC with 'A++' Grade (Cycle III) &

Accredited by NBA (B.E - CSE, EEE, ECE, Mech & B.Tech.IT)

COIMBATORE-641 035, TAMIL NADU



$$f(x) = \frac{x^2}{3} + \sum_{n=1}^{\infty} \frac{4x^2 (-1)^n}{n^2 \pi^2} \cos n\pi$$

$$= \frac{x^2}{3} + \frac{4x^2}{\pi^2} \sum_{n=1}^{\infty} \frac{(-1)^n (-1)^n}{n^2}$$

$$f(x) = \frac{x^2}{3} + \frac{4x^2}{\pi^2} \sum_{n=1}^{\infty} \frac{1}{n^2} \quad \text{--- (3)}$$

$$\therefore (-1)^n (-1)^n = 1$$

To find  $f(x)$ :

$$f(x) = x^2 \text{ in } -1 < x < 1$$

$x=1$  is end point of  $(-1, 1)$

$$f(1) = 1^2$$

$$f(-1) = (-1)^2 = 1^2$$

$$f(1) = f(-1) = 1^2$$

Here  $x=1$  is a point of continuity

$$\therefore f(1) = 1^2$$

Substitute  $f(x) = 1^2$  in (3), we get

$$1^2 = \frac{1^2}{3} + \frac{4 \cdot 1^2}{\pi^2} \sum_{n=1}^{\infty} \frac{1}{n^2}$$

$$\frac{4 \cdot 1^2}{\pi^2} \sum_{n=1}^{\infty} \frac{1}{n^2} = 1^2 - \frac{1^2}{3}$$

$$= \frac{2 \cdot 1^2}{3}$$



# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Approved by AICTE, New Delhi, Affiliated to Anna University, Chennai

Accredited by NAAC-UGC with 'A++' Grade (Cycle III) &

Accredited by NBA (B.E - CSE, EEE, ECE, Mech & B.Tech.IT)

COIMBATORE-641 035, TAMIL NADU



$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{2l^2}{3} \frac{\pi^2}{4l^2}$$

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}$$

$$\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots = \frac{\pi^2}{6}$$

(ii) To deduce  $\frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \dots = \frac{\pi^2}{12}$

Put  $x=0$  in (2) we get

$$f(0) = \frac{1^2}{3} + \frac{4l^2}{\pi^2} \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2} \quad \text{--- (4)}$$

To find  $f(0)$ :

$$f(x) = x^2$$

$\therefore x=0$  lies inside  $(-l, l)$

Here  $x=0$  is a point of continuity

$$\therefore f(0) = 0$$

Substitute  $f(0) = 0$  in (4) we get

$$0 = \frac{1^2}{3} + \frac{4l^2}{\pi^2} \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2}$$

$$\frac{4l^2}{\pi^2} \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2} = -\frac{1^2}{3}$$

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{n^2} = -\frac{1^2}{3} \frac{\pi^2}{4l^2}$$



# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Approved by AICTE, New Delhi, Affiliated to Anna University, Chennai

Accredited by NAAC-UGC with 'A++' Grade (Cycle III) &

Accredited by NBA (B.E - CSE, EEE, ECE, Mech & B.Tech.IT)

COIMBATORE-641 035, TAMIL NADU



$$(-1) \frac{1}{1^2} + \frac{1}{2^2} + (-1) \frac{1}{3^2} + \dots = \frac{-\pi^2}{12}$$

~~$\frac{1}{1^2}$~~

Ans:  $\boxed{\frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \dots = \frac{\pi^2}{12}}$

② Find the Fourier series  $f(x) = |x|$ ,  $-\pi < x < \pi$  and deduce that  $\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots = \frac{\pi^2}{8}$

Solution:

$f(x)$  is defined in  $(-\pi, \pi)$

In  $(-\pi, \pi)$  check whether  $f(x)$  is even or odd

$$f(x) = |x|$$

$$f(-x) = |-x| = |x| = f(x)$$

$$f(-x) = f(x)$$

$\therefore f(x)$  is an even function

$$\therefore b_n = 0$$

The Fourier series for  $f(x)$  in  $(-\pi, \pi)$  is given by.

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos nx \quad \text{--- ①}$$

$$a_0 = \frac{2}{\pi} \int_0^{\pi} f(x) dx$$



# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Approved by AICTE, New Delhi, Affiliated to Anna University, Chennai

Accredited by NAAC-UGC with 'A++' Grade (Cycle III) &

Accredited by NBA (B.E - CSE, EEE, ECE, Mech & B.Tech.IT)

COIMBATORE-641 035, TAMIL NADU



$$= \frac{2}{\pi} \int_0^{\pi} x \, dx$$

$$\therefore |x| = \begin{cases} -x, & -\pi < x < 0 \\ x, & 0 < x < \pi \end{cases}$$

$$= \frac{2}{\pi} \left[ \frac{x^2}{2} \right]_0^{\pi}$$

$$= \frac{2}{\pi} \times \frac{\pi^2}{2}$$

$$a_0 = \pi$$

$$a_n = \frac{2}{\pi} \int_0^{\pi} f(x) \cos nx \, dx$$

$$= \frac{2}{\pi} \int_0^{\pi} x \cos nx \, dx$$

$$= \frac{2}{\pi} \left[ x \left( \frac{\sin nx}{n} \right) - \left( \frac{\cos nx}{n^2} \right) \right]_0^{\pi}$$

$$= \frac{2}{\pi n^2} [\cos nx]_0^{\pi}$$

$$= \frac{2}{\pi n^2} [(-1)^n - 1]$$

$$a_n = \begin{cases} -\frac{4}{\pi n^2} & ; n = 1, 3, 5, \dots \\ 0 & ; n = 2, 4, 6, \dots \end{cases}$$

Substituting the values of  $a_0, a_n$  in (1), we get.





# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Approved by AICTE, New Delhi, Affiliated to Anna University, Chennai

Accredited by NAAC-UGC with 'A++' Grade (Cycle III) &

Accredited by NBA (B.E - CSE, EEE, ECE, Mech & B.Tech.IT)

COIMBATORE-641 035, TAMIL NADU



$$f(x) = \frac{\pi}{2} + \sum_{n=1,3}^{\infty} \frac{-4}{\pi n^2} \cos nx \quad \text{--- (2)}$$

i) To deduce  $\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots = \frac{\pi^2}{8}$

Put  $x=0$  in (2), we get

$$\begin{aligned} f(0) &= \frac{\pi}{2} + \sum_{n=1,3}^{\infty} \frac{-4}{\pi n^2} \\ &= \frac{\pi}{2} - \frac{4}{\pi} \sum_{n=1}^{\infty} \frac{1}{n^2} \quad \text{--- (3)} \end{aligned}$$

To find  $f(0)$ :

$x=0$  lies inside  $(-\pi, \pi)$ .

Here  $x=0$  is a point of continuity

$$f(x) = |x|$$

$$f(0) = |0| = 0$$

Substitute  $f(0)=0$  in (3), we get

$$0 = \frac{\pi}{2} - \frac{4}{\pi} \sum_{n=1,3}^{\infty} \frac{1}{n^2}$$

$$-\frac{4}{\pi} \sum_{n=1,3}^{\infty} \frac{1}{n^2} = -\frac{\pi}{2}$$

$$\sum_{n=1,3}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{8}$$

$$\boxed{\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots = \frac{\pi^2}{8}}$$

Hence proved