



## Applications of Laplace transforms to

### Differential Equations:

If  $L[f(t)] = F(s)$  then

$$L[y'(t)] = sL(y) - y(0)$$

$$L[y''(t)] = s^2L(y) - sy(0) - y'(0)$$

- ① Solve the differential equations using LT  
 $y'' + 4y' + 4y = e^{-t}$  given that  $y(0) = 0$  and  $y'(0) = 0$ .

Soln:  $y'' + 4y' + 4y = e^{-t}$

Taking LT on both sides,

$$L(y'' + 4y' + 4y) = L(e^{-t})$$

$$L(y'') + 4L(y') + 4L(y) = \frac{1}{s+1}$$

$$[s^2L(y) - sy(0) - y'(0)] + 4[sL(y) - y(0)]$$

$$+ 4L(y) = \frac{1}{s+1}$$



Given:  $y(0) = 0, y'(0) = 0$

$$\Rightarrow [s^2 L(y) - s y(0) - y'(0)] + 4 [s L(y) - y(0)] + 4 L(y) = \frac{1}{s+1}$$

$$\Rightarrow s^2 L(y) + 4s L(y) + 4 L(y) = \frac{1}{s+1}$$

$$\Rightarrow (s^2 + 4s + 4) L(y) = \frac{1}{s+1}$$

$$\Rightarrow L(y) (s+2)^2 = \frac{1}{s+1}$$

$$\Rightarrow L(y) = \frac{1}{(s+1)(s+2)^2}$$

$$y = L^{-1} \left[ \frac{1}{(s+1)(s+2)^2} \right]$$

$$\frac{1}{(s+1)(s+2)^2} = \frac{A}{s+1} + \frac{B}{s+2} + \frac{C}{(s+2)^2}$$

$$1 = A(s+2)^2 + B(s+2)(s+1) + C(s+1)$$

Put  $s = -2 \Rightarrow C = -1$

Put  $s = -1 \Rightarrow A = 1$

Put  $s = 0 \Rightarrow B = -1$

$$\frac{1}{(s+1)(s+2)^2} = \frac{1}{s+1} - \frac{1}{s+2} - \frac{1}{(s+2)^2}$$

$$y = L^{-1} \left( \frac{1}{(s+1)(s+2)^2} \right)$$

$$= L^{-1} \left( \frac{1}{s+1} \right) - L^{-1} \left( \frac{1}{s+2} \right) - L^{-1} \left( \frac{1}{(s+2)^2} \right)$$

$$= e^{-t} - e^{-2t} - t e^{-2t}$$

$$= e^{-t} - e^{-2t} - t e^{-2t}$$