



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



Unit 3-Differential Calculus

Evolutes

Q. Find the eqn. of evolute of the astroid

$$x^{2/3} + y^{2/3} = a^{2/3}$$

Soln.

Parametric form

$$\begin{aligned} x &= a \cos^3 \theta & y &= a \sin^3 \theta \\ \frac{dx}{d\theta} &= 3a \cos^2 \theta (-\sin \theta) & \frac{dy}{d\theta} &= 3a \sin^2 \theta (\cos \theta) \end{aligned}$$

$$\begin{aligned} \frac{dy}{dx} &= \frac{dy}{d\theta} \times \frac{d\theta}{dx} = \frac{3a \sin^2 \theta \cos \theta}{-3a \cos^2 \theta \sin \theta} \\ &= -\frac{\sin \theta}{\cos \theta} \end{aligned}$$

$$y_1 = -\tan \theta$$

$$\begin{aligned} \frac{d^2y}{dx^2} &= \frac{d}{d\theta} \left[\frac{dy}{dx} \right] \frac{d\theta}{dx} \\ &= \frac{d}{d\theta} [-\tan \theta] \cdot \frac{1}{-3a \cos^2 \theta \sin \theta} \\ &= \frac{-\sec^2 \theta}{-3a \cos^2 \theta \sin \theta} \end{aligned}$$

$$y_2 = \frac{\sec^4 \theta}{3a \sin \theta} = \frac{1}{3a} \csc \theta \sec^4 \theta$$

$$\begin{aligned} x &= x - \frac{y_1(1+y_1^2)}{y_2} \\ &= a \cos^3 \theta - \frac{(-\tan \theta)(1+\tan^2 \theta)}{\frac{1}{3a} \csc \theta \sec^4 \theta} \\ &= a \cos^3 \theta + 3a \frac{\sin \theta}{\cos \theta} \sin \theta \cos^4 \theta \sec^2 \theta \\ &= a \cos^3 \theta + 3a \sin^2 \theta \cos^3 \theta \frac{1}{\cos^2 \theta} \end{aligned}$$

Scanned with CamScanner



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



Unit 3-Differential Calculus

Evolutes

$$\begin{aligned}
 \bar{x} &= a \cos^3 \theta + 3a \sin^2 \theta \cos \theta \rightarrow (1) \\
 \bar{y} &= y + \frac{1+y_1^2}{y_2} \\
 &= a \sin^3 \theta + \frac{1+\tan^2 \theta}{\frac{1}{2a} \csc \theta \sec^4 \theta} \\
 &= a \sin^3 \theta + 3a \frac{\sec^2 \theta}{\csc \theta \sec^4 \theta} \\
 &= a \sin^3 \theta + 3a \frac{1}{\csc \theta \sec^2 \theta} \\
 \bar{y} &= a \sin^3 \theta + 3a \sin \theta \cos^2 \theta \rightarrow (2) \\
 (1) + (2) &\Rightarrow \bar{x} + \bar{y} = a \cos^3 \theta + 3a \sin^2 \theta \cos \theta + a \sin^3 \theta + \\
 &\quad 3a \sin \theta \cos^2 \theta \\
 &= a [\cos^3 \theta + 3\sin^2 \theta \cos \theta + 3\sin \theta \cos^2 \theta + \sin^3 \theta] \\
 \bar{x} + \bar{y} &= a [\cos \theta + \sin \theta]^3 \rightarrow (3) \\
 (1) - (2) &\Rightarrow \bar{x} - \bar{y} = a \cos^3 \theta + 3a \sin^2 \theta \cos \theta - a \sin^3 \theta - 3a \sin \theta \cos^2 \theta \\
 \bar{x} - \bar{y} &= a [\cos \theta - \sin \theta]^3 \rightarrow (4) \\
 \text{Taking power as } 2/3 \text{ on both sides of (1) & (2)} & \\
 (\bar{x} + \bar{y})^{2/3} &= a^{2/3} [\cos \theta + \sin \theta]^{3 \times 2/3} \\
 (\bar{x} + \bar{y})^{2/3} &= a^{2/3} [\cos \theta + \sin \theta]^2 \rightarrow (5) \\
 \text{and } (\bar{x} - \bar{y})^{2/3} &= a^{2/3} [\cos \theta - \sin \theta]^2 \rightarrow (6) \\
 (5) + (6) &\Rightarrow (\bar{x} + \bar{y})^{2/3} + (\bar{x} - \bar{y})^{2/3} = a^{2/3} [\cos \theta + \sin \theta]^2 + \\
 &\quad a^{2/3} [\cos \theta - \sin \theta]^2 \\
 &= a^{2/3} [\cos^2 \theta + \sin^2 \theta + 2 \cos \theta \sin \theta + \cos^2 \theta + \sin^2 \theta \\
 &\quad - 2 \sin \theta \cos \theta] \\
 &= a^{2/3} [1 + 1] \\
 (\bar{x} + \bar{y})^{2/3} + (\bar{x} - \bar{y})^{2/3} &= 2a^{2/3}
 \end{aligned}$$

Scanned with CamScanner



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



Unit 3-Differential Calculus

Evolutes

Unit-3 continuation...

Q. Find the evolute of $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$

Soln.

$$\begin{aligned} \text{Take } x &= a \sec \theta & y &= b \tan \theta \\ \frac{dx}{d\theta} &= a \sec \theta \tan \theta & \frac{dy}{d\theta} &= b \sec^2 \theta \end{aligned}$$

$$\text{Now } \frac{dy}{dx} = \frac{dy}{d\theta} \cdot \frac{d\theta}{dx}$$

$$= b \sec^2 \theta$$

$$\frac{dy}{dx} = \frac{b}{a} \frac{\sec \theta}{\tan \theta}$$

$$\frac{d^2y}{dx^2} = \frac{d}{d\theta} \left[\frac{dy}{dx} \right] \frac{d\theta}{dx}$$

$$= \frac{d}{d\theta} \left[\frac{b}{a} \frac{\sec \theta}{\tan \theta} \right] \frac{1}{a \sec \theta \tan \theta}$$

$$= \frac{b}{a} \left[\frac{\tan \theta \sec \theta \tan \theta - \sec \theta \sec^2 \theta}{\tan^2 \theta} \right] \frac{1}{a \sec \theta \tan \theta}$$

$$= \frac{b}{a} \left[\frac{\tan^2 \theta \sec \theta - \sec^3 \theta}{\tan^2 \theta} \right] \frac{1}{a \sec \theta \tan \theta}$$

$$= \frac{b}{a} \left[\frac{\tan^2 \theta - \sec^2 \theta}{\tan^2 \theta} \right] \sec \theta \frac{1}{a \sec \theta \tan \theta}$$

$$= \frac{b}{a} \frac{-1}{\tan^2 \theta} \frac{1}{a \tan \theta}$$

$$\frac{d^2y}{dx^2} = -\frac{b}{a^2 \tan^3 \theta}$$

$$\bar{x} = x - \frac{y_1 [1 + y_1^2]}{y_2} = a \sec \theta - \frac{\frac{b}{a} \frac{\sec \theta}{\tan \theta} \left[1 + \frac{b^2 \sec^2 \theta}{a^2 \tan^2 \theta} \right]}{-\frac{b}{a^2 \tan^3 \theta}}$$



Scanned with CamScanner



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



Unit 3-Differential Calculus

Evolutes

Exam-N

$$\begin{aligned}
 &= a \sec \theta + \frac{b}{a} \frac{\sec \theta}{\tan \theta} \quad \frac{a^2 \tan^2 \theta + b^2 \sec^2 \theta}{a^2 + \tan^2 \theta} \times \frac{a^2 \tan^3 \theta}{b} \\
 &= a \sec \theta + \frac{a^2}{a} \sec \theta \tan^2 \theta + \frac{b^2}{a} \sec \theta \sec^2 \theta \\
 &= a \sec \theta + a \sec \theta \tan^2 \theta + \frac{b^2}{a} \sec^3 \theta \\
 &= a \sec \theta [1 + \tan^2 \theta] + \frac{b^2}{a} \sec^3 \theta \\
 &= a \sec \theta [\sec^2 \theta] + \frac{b^2}{a} \sec^3 \theta = a \sec^3 \theta + \frac{b^2}{a} \sec^3 \theta
 \end{aligned}$$

$$\begin{aligned}
 \bar{x} &= \left[\frac{a^2 + b^2}{a} \right] \sec^3 \theta \\
 \bar{y} &= b \tan \theta + \frac{\left[1 + \frac{b^2 \sec^2 \theta}{a^2 \tan^2 \theta} \right]}{-\frac{b}{a^2} \tan^3 \theta} \\
 &= b \tan \theta - \frac{a^2}{b} \tan^3 \theta \left[\frac{a^2 \tan^2 \theta + b^2 \sec^2 \theta}{a^2 \tan^2 \theta} \right] \\
 &= b \tan \theta - \frac{\tan \theta}{b} [a^2 \tan^3 \theta + b^2 \sec^2 \theta] \\
 &= b \tan \theta - \frac{a^2}{b} \tan^3 \theta - b \sec^2 \theta \tan \theta \\
 &= b \tan \theta [1 - \sec^2 \theta] - \frac{a^2}{b} \tan^3 \theta \\
 &= -b \tan \theta \tan^2 \theta - \frac{a^2}{b} \tan^3 \theta = -b \tan^3 \theta - \frac{a^2}{b} \tan^3 \theta \\
 &= \left[b - \frac{a^2}{b} \right] \tan^3 \theta \\
 &= \left[\frac{-b^2 - a^2}{b} \right] \tan^3 \theta \\
 \bar{y} &= -\left[\frac{a^2 + b^2}{b} \right] \tan^3 \theta
 \end{aligned}$$

$$\begin{aligned}
 \text{Now, } \bar{x} &= \left[\frac{a^2 + b^2}{a} \right] \sec^3 \theta & \bar{y} &= -\left[\frac{a^2 + b^2}{b} \right] \tan^3 \theta \\
 \Rightarrow \sec^3 \theta &= \frac{a \bar{x}}{a^2 + b^2} & \Rightarrow \tan^3 \theta &= \frac{-b \bar{y}}{(a^2 + b^2)} \\
 \Rightarrow \sec^2 \theta &= \left[\frac{a \bar{x}}{a^2 + b^2} \right]^{2/3} & \Rightarrow \tan^2 \theta &= \left[\frac{b \bar{y}}{a^2 + b^2} \right]^{2/3} \\
 &\Downarrow (1) & &\Downarrow (2)
 \end{aligned}$$



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



Unit 3-Differential Calculus

Evolutes

NOW (1) - (2),

$$\sec^2 \theta - \tan^2 \theta = \left[\frac{a\bar{x}}{a^2+b^2} \right]^{2/3} - \left[\frac{b\bar{y}}{a^2+b^2} \right]^{2/3}$$

$$\Rightarrow (a^2+b^2)^{2/3} = (\bar{x})^{2/3} - (\bar{y})^{2/3}$$

Replace \bar{x} by x and \bar{y} by y .

$$\Rightarrow (ax)^{2/3} - (by)^{2/3} = (a^2+b^2)^{2/3}$$

Show that the evolute of the rectangular hyperbola $xy=c^2$ is $(x+y)^{2/3} - (x-y)^{2/3} = (4c)^{2/3}$

Soln.

Take $x = ct$ and $y = \frac{c}{t}$

$$\frac{dx}{dt} = c \quad \frac{dy}{dt} = -\frac{c}{t^2}$$

$$\frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dt}{dx} = -\frac{c}{t^2} \times \frac{1}{c} = -\frac{1}{t^2}$$

$$\frac{d^2y}{dx^2} = \frac{2}{ct^3}$$

$$\bar{x} = 3ct + \frac{\frac{1}{t^2}[1 + \frac{1}{t^4}]}{\frac{2}{ct^3}} = ct + \frac{1}{t^2} \left[\frac{t^4+1}{t^4} \right] \frac{ct^3}{2}$$

$$= ct + \frac{c}{2t^3}[t^4+1]$$

$$= ct + \frac{ct}{2} + \frac{c}{2t^3}$$

$$\bar{x} = \frac{3ct}{2} + \frac{c}{2t^3}$$

$$\bar{y} = \frac{c}{t} + \frac{\frac{1}{t^2}[1 + \frac{1}{t^4}]}{\frac{2}{ct^3}} = \frac{c}{t} + \frac{ct^3}{2} \left[\frac{t^4+1}{t^4} \right]$$

$$= \frac{c}{t} + \frac{c}{2t} [t^4+1] = \frac{c}{t} + \frac{ct^3}{2} + \frac{c}{2t}$$

$$\bar{y} = \frac{3c}{2t} + \frac{ct^3}{2}$$

Scanned with CamScanner



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



Unit 3-Differential Calculus

Evolutes

$$\text{Now, } \bar{x} + \bar{y} = \frac{3Cz}{2} + \frac{C}{2z^3} + \frac{3C}{2z} + \frac{Cz^3}{2}$$

$$= \frac{C}{2} [3z + \frac{1}{z^3} + \frac{3}{z} + z^3]$$

$$\bar{x} + \bar{y} = \frac{C}{2} [z + \frac{1}{z}]^3 \rightarrow (1)$$

$$\text{and } \bar{x} - \bar{y} = \frac{3Cz}{2} + \frac{C}{2z^3} - \frac{3C}{2z} - \frac{Cz^3}{2}$$

$$= \frac{C}{2} [3z + \frac{1}{z^3} - \frac{3}{z} - z^3]$$

$$\bar{x} - \bar{y} = -\frac{C}{2} [z - \frac{1}{z}]^3 \rightarrow (2)$$

Taking power as $2/3$ on both sides of (1) & (2)

$$(\bar{x} + \bar{y})^{2/3} = \left(\frac{C}{2}\right)^{2/3} [z + \frac{1}{z}]^2$$

$$(\bar{x} - \bar{y})^{2/3} = \left(\frac{C}{2}\right)^{2/3} [z - \frac{1}{z}]^2$$

$$\text{Now, } (\bar{x} + \bar{y})^{2/3} - (\bar{x} - \bar{y})^{2/3}$$

$$= \left(\frac{C}{2}\right)^{2/3} [(z + \frac{1}{z})^2 - (z - \frac{1}{z})^2]$$

$$= \left(\frac{C}{2}\right)^{2/3} (4)$$

Replace \bar{x} by x & \bar{y} by y ,

$$(x + y)^{2/3} - (x - y)^{2/3} = 4 \left(\frac{C}{2}\right)^{2/3} = 2^2 2^{-2/3} C^{2/3}$$

$$= 2^{4/3} C^{2/3}$$

$$= (4C)^{2/3} \quad (2^{4/3} = 4^{2/3})$$



Unit 3-Differential Calculus

Evolutes

Q5.

Show that the evolute of cycloid $x = a(t + \sin t)$, $y = a(1 - \cos t)$ is $x = a(t - \sin t)$; $y - 2a = a(1 + \cos t)$

Soln.

Given $x = a(t + \sin t)$

$$\frac{dx}{dt} = a(1 + \cos t)$$

$$\Rightarrow \frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dt}{dx}$$

$$= a \sin t \times \frac{1}{a(1 + \cos t)}$$

$$= \frac{\sin t}{1 + \cos t}$$

$$= \frac{2 \sin t/2 \cos t/2}{2 \cos^2 t/2}$$

$$= \tan t/2$$

$$\frac{d^2y}{dx^2} = \frac{d}{dt} \left(\frac{dy}{dx} \right) \frac{dt}{dx}$$

$$= \frac{d}{dt} \left[\tan \frac{t}{2} \right] \frac{1}{a(1 + \cos t)}$$

$$= \sec^2 \frac{t}{2} \left(\frac{1}{2} \right) \frac{1}{a(1 + \cos t)}$$

$$= \frac{\sec^2 t/2}{2a [2 \cos^2 t/2]}$$

$$= \frac{\sec^4 t/2}{4a}$$

$$\bar{x} = x - \frac{y_1 (1 + y_1^2)}{y_2} = a(t + \sin t) - \frac{\tan t/2 [1 + \tan^2 t/2]}{\sec^4 t/2}$$

$$= a(t + \sin t) - \frac{4 \tan t/2 [\sec^2 t/2]}{\sec^4 t/2}$$

Scanned with CamScanner



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



Unit 3-Differential Calculus

Evolutes

$$\begin{aligned}
 &= a(t + \sin t) - 4a \sin t/2 \cos^2 t/2 \\
 &= at + a \sin t - 2a [2 \sin t/2 \cos^2 t/2] \\
 &= at + a \sin t - 2a \sin t \cos^2 t/2 \\
 &= at + a \sin t + 2a \sin t \\
 &= at - a \sin t \\
 \bar{x} &= a(t - \sin t) \rightarrow (1) \\
 \bar{y} &= y + \frac{1 + y_1^2}{y_2} \\
 &= a(1 - \cos t) + \frac{1 + \tan^2 t/2}{\sec^4 t/2} \\
 &= a(1 - \cos t) + \frac{4a \sec^2 t/2}{\sec^4 t/2} \\
 &= a(1 - \cos t) + 4a \cos^2 t/2 \\
 &= a - a \cos t + 2a [\cos^2 t/2] \\
 &= a - a \cos t + 2a[1 + \cos t] \quad \because \frac{1 + \cos 2\theta}{2} = \cos^2 \theta \\
 &= a - a \cos t + 2a + 2a \cos t \\
 \bar{y} - 2a &= a + a \cos t \rightarrow (2) \\
 \text{Replace } \bar{x} \text{ by } x \text{ & } \bar{y} \text{ by } y \text{ in (1) & (2),} \\
 x &= a(t - \sin t) \text{ &} \\
 (y - 2a) &= a + a \cos t = a(1 + \cos t)
 \end{aligned}$$

Ques

- J. Find the eqn. of evolute $x^{2/3} + y^{2/3} = a^{2/3}$
- Q. Show that the evolute of the cycloid $x = a(\theta - \sin \theta); y = a(1 - \cos \theta)$ is another cycloid $x = a(\theta + \sin \theta); y = -a(1 - \cos \theta)$



Scanned with CamScanner