



UNIT-3

Free Electron theory of metals

Classical free electron theory is based on the following postulates:

1. A solid metal is composed of atoms and the atoms have nucleus, around which there are revolving electrons.
2. In a metal the valance electrons of atoms are free to move throughout the volume of the metal like gas molecules of a perfect gas in a container
3. The free electrons move in a random directions and collide with either positive ions fixed to the lattice or other free electrons and collisions are elastic in nature i.e. there is no loss of energy.
4. The movement of free electrons obeys the classical kinetic theory of gasses. The mean K.E. of a free electron is equal to that of gas molecule $\frac{3}{2} kT$
5. The electron velocities in a metal obey Maxwell-Boltzman distribution of velocities.
6. The free electrons move in a uniform potential field due to ions fixed in the lattice
7. When an electric field is applied to the metal the free electrons are accelerated. The accelerated electrons move in opposite direction of the applied.
8. The electric conduction is due to the free electrons only.

ROOT MEAN SQUARE (R.M.S.) VELOCITY:

Let \bar{c} be the r.m.s velocity of the free electron. then the

$$\text{Kinetic energy} = \frac{1}{2} m \bar{c}^2$$

But according to the classical free electron theory the mean

$$\text{Kinetic Energy} = \frac{3}{2} kT$$

$$\frac{1}{2} m \bar{c}^2 = \frac{3}{2} kT$$

$$\Rightarrow \bar{c} = \sqrt{\frac{3KT}{m}} \quad \text{where } \bar{c} = \text{root mean square velocity}$$



MEAN FREE PATH AND MEAN COLLISION TIME

The average distance travelled by an electron between two successive collisions in the presence of applied field is known as 'Mean free path'.


The time taken by an electron between two successive collisions is known as "Mean Collision Time" of the electron

$$\lambda = \sqrt{\frac{m}{3KT}}$$

DRIFT VELOCITY (v_d):

It is the average velocity acquired by the free electrons of a metal in a particular direction during the application of the electric field.

ELECTRICAL CONDUCTIVITY IN METALS:



Let us consider a conductor of length l and area of cross section A

The volume of the conductor Al

If there are n number of electrons per unit volume of the

metal then the total number of electrons in the metal

$$Aln$$

If e is the charge of the electron then the total charge q due to all electrons in the conductor is given by $q = Aln.e$

Let t be the time taken by the electron to move from one end to other end then

$$\text{Current (I)} = \frac{\text{charge}}{\text{time}} = \frac{q}{t} = \frac{Aln.e}{t}$$



SNS COLLEGE OF TECHNOLOGY

But $I = n e v_d A$ (An Autonomous Institution)



$$\Rightarrow v_d = \frac{I}{n e A}$$

Where J current density $= \frac{I}{A}$

In a metal the current density J is given by the equation

$$J = n e v_d \dots \dots \dots (1)$$

Where n number of electrons per Unit volume, e electron charge and v_d drift velocity. If E is the applied electric field then the electric force acting on a free electron is given by

$$F = eE \dots \dots \dots (2)$$

From Newton's IInd law $F = ma \dots \dots \dots (3)$

From (2) and (3) $ma = eE$

i.e. $a = \frac{eE}{m}$

but a drift velocity/collision time $\frac{v_d}{\tau}$

$$v_d = a \tau = \frac{eE \tau}{m}$$

$$J = n e \frac{eE \tau}{m} = \frac{n e^2 E \tau}{m} \dots \dots \dots (4)$$



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)



But from microscopic form of ohms law

$$J = \sigma E \dots \dots \dots (5)$$

On comparing Eq(4)&(5)

$$\text{Conductivity } \sigma = \frac{ne^2 \tau}{m} \text{ or Resistivity } \rho = \frac{m}{ne^2 \tau}$$

Conductivity may also be expressed in terms of mobility (μ) which is defined as drift velocity per unit electric field

$$\mu = \frac{v_d}{E} = \frac{e \tau}{m}$$

From (4) $\sigma = ne \mu$

RELAXATION TIME (τ)

Under the influence of an external electric field free electrons attain a directional velocity of motion. If the field is switched off the velocity starts decreasing exponentially. Such a process that tends to restore equilibrium is called relaxation process.

If v_0 is the velocity at $t = 0$ at which the field is switched

off. The velocity at any time is given by

$$v = v_0 e^{-t/\tau}$$

In the above expression τ = relaxation

time If $t = \tau$

$$v = v_0 e^{-1} = \frac{v_0}{e}$$

Relaxation time τ is defined as the time required for the electron to reduce its velocity to $\frac{1}{e}$ of

its initial value. (OR) time taken for the drift velocity to decay to $\frac{1}{e}$ of its initial value.



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

