



(An Autonomous Institution) Coimbatore 35

### DEPARTMENT OF MATHEMATICS

UNIT-I TESTING OF HYPOTHESIS

CHI SQUARE TEST FOR INDEPENDENCE OF ATTRIBUTES.

I) One the basis of information noted below, find out whether the new treatment & compeasitively superior to the conventional one.

	Favourable.	Not Favourable	weal	
New	60	30	90	
Conventional	40	70	110	
n: total	100	100 ->	200	

	90 x 100 : 45	90 x 100 : 45	
1	110 × 100 : 55	110 x 100 , 55	, Purary or bire
	200	200	DIT- LA





(An Autonomous Institution)
Coimbatore— 35

### **DEPARTMENT OF MATHEMATICS**

#### UNIT-I TESTING OF HYPOTHESIS

00 Ei 0i-Ei (0i-Ei)/Ei

60 45 15 5

30 45 -15 5

40 55 -15 4.09

40 55 15 4.09

$$\sum_{EL} (a:-E_L)^2 \cdot 18.18$$

step 1: Formulating Ho & H .:

Ho: There is no difference between mew & conventioned breatment.

HI: There is difference hetween mew &

conventinal treatment.

step 2. Los al x = 5%.

slep 3: Test statistics,  $\chi^2 = \frac{5(0:-5:)^2}{E:}$ = 18.18

step 4: Degrees & Freedom, N = ((3-1) + (+-1))

V= (2-1 \* 2-1)

= 1 \*1

. . tab value, Xx = 3.841

top 5 : Conclusion:

X2 = 18.18 > 3.841 = X2

. Ho is rejected at 5 1/. Los

treatment.





(An Autonomous Institution)
Coimbatore— 35

## **DEPARTMENT OF MATHEMATICS**

## **UNIT-I TESTING OF HYPOTHESIS**

2) Two resease	cheis A	and B ac	lopted differen	+ techniques
1.1 . 1.9	He strid	ente leve	l, can you se	ey meet inc
1-1-80,1100 00	lunteel h	y them an	e ngrufican	F ?
1	polous	ava. A	vg. Above All	g. Genius 101d
A	40	3	3	
		1.7	44	10 200
В	101	5 N	60	12 300
-to-lal	126	9.5	69	
din				s and
70 be				
	100 x 126: 4	2 100 × 93	31 100 x 69 2	3 300 4
	200 x 126 8	300 x 93	: 62 200x 69	8. 500x13
0;	Eį	Oc - E;	(0;-Ei) /E:	
AD	42	- 2	0.0952	
33	31	. 2	0-129	
35	23	2	0.173	
2	4	- 2	1	
86	84	2	0.047	
60	62	- 2	0.064	
44	46	- 2	0.086	
16	8	Ş	0.5	in the
		2 (0i	Ec)2. 2.094	





(An Autonomous Institution) Coimbatore 35

### DEPARTMENT OF MATHEMATICS

UNIT-I TESTING OF HYPOTHESIS

Step 1: Formulating Ho and Hi:

Ho: Fihere is no difference tretivosen the mo,

H.: There is difference between the two remarcher

(step 2 : Los at x = 5%.

step 3: Test statuties,  $\chi^2 = \underbrace{\mathbb{E}\left(0i-\epsilon_i\right)^2}_{Ei}$ 

step 4: peyers of freedom, v= ((4-1) \* (2-1))

= (3 \* 1) ..

: Tab value is 2 = 4.115

glap 5: Conclusion.

82 = 2.097 <11115 = 22x

: Ho is accepted at 5% Los

(a) There is no difference between The two severaschers,