



DEPARTMENT OF MATHEMATICS

14) Find the values of a and b so that the surface $ax^3 - by^2z = (a+3)x^2$ and $4x^2y - z^3 = 11$ may cut orthogonally at $(2, -1, -3)$.

Soln:

$$a = -7/3 \quad \& \quad b = 64/9$$

DIVERGENCE OF A VECTOR POINT FUNCTION:

Let \vec{F} be any given continuously differentiable vector point function then the divergence of \vec{F} is defined as,

$$\begin{aligned} \text{div } \vec{F} &= \nabla \cdot \vec{F} = \left(\vec{i} \frac{\partial}{\partial x} + \vec{j} \frac{\partial}{\partial y} + \vec{k} \frac{\partial}{\partial z} \right) \cdot \vec{F} \\ &= \vec{i} \cdot \frac{\partial \vec{F}}{\partial x} + \vec{j} \cdot \frac{\partial \vec{F}}{\partial y} + \vec{k} \cdot \frac{\partial \vec{F}}{\partial z} \end{aligned}$$

Note :

- $\nabla \cdot \vec{F}$ is a scalar point function.
- If $\vec{F} = F_1 \vec{i} + F_2 \vec{j} + F_3 \vec{k}$ be a continuously differentiable vector point function then,

$$\text{div } \vec{F} = \frac{\partial F_1}{\partial x} + \frac{\partial F_2}{\partial y} + \frac{\partial F_3}{\partial z}$$

Solenoidal vector:

A vector \vec{F} is said to be solenoidal vector if $\text{div } \vec{F} = 0$.

CURL OF A VECTOR POINT FUNCTION:

Let $\vec{F} = F_1 \vec{i} + F_2 \vec{j} + F_3 \vec{k}$ be any given continuously differentiable vector point function, the curl or rotation of \vec{F} is defined as,



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Approved by AICTE, New Delhi, Affiliated to Anna University, Chennai

Accredited by NAAC-UGC with 'A++' Grade (Cycle III) &

Accredited by NBA (B.E - CSE, EEE, ECE, Mech & B.Tech.IT)

COIMBATORE-641 035, TAMIL NADU

DEPARTMENT OF MATHEMATICS

$$\begin{aligned}\text{curl } \vec{F} &= \nabla \times \vec{F} \\ &= \left(\vec{i} \frac{\partial}{\partial x} + \vec{j} \frac{\partial}{\partial y} + \vec{k} \frac{\partial}{\partial z} \right) \times \vec{F} \\ &= \vec{i} \times \frac{\partial \vec{F}}{\partial x} + \vec{j} \times \frac{\partial \vec{F}}{\partial y} + \vec{k} \times \frac{\partial \vec{F}}{\partial z} \\ &= \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ F_1 & F_2 & F_3 \end{vmatrix}\end{aligned}$$

Note: $\nabla \times \vec{F}$ is a vector point function.

IRROTATIONAL VECTOR:

A vector \vec{F} is said to be irrotational if

$$\nabla \times \vec{F} = 0$$

$$\text{i.e., } \text{curl } \vec{F} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ F_1 & F_2 & F_3 \end{vmatrix} = 0$$

CONSERVATIVE VECTOR FIELD:

If a vector point function \vec{F} is expressible as the gradient of a scalar point function ϕ , then

\vec{F} is conservative i.e., \vec{F} is conservative if $\vec{F} = \nabla \phi$.

Here ϕ is called scalar potential.

\vec{F} is conservative force if $\text{curl } \vec{F} = 0$.



DEPARTMENT OF MATHEMATICS

PROBLEMS :

① Prove that $\text{curl}(\nabla\phi) = 0$ (or) $\nabla \times \nabla\phi = 0$.

Soln:

$$\nabla = \vec{i} \frac{\partial}{\partial x} + \vec{j} \frac{\partial}{\partial y} + \vec{k} \frac{\partial}{\partial z}$$

$$\nabla\phi = \vec{i} \frac{\partial\phi}{\partial x} + \vec{j} \frac{\partial\phi}{\partial y} + \vec{k} \frac{\partial\phi}{\partial z}$$

$$\text{Curl}(\nabla\phi) = \nabla \times \nabla\phi = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ \frac{\partial\phi}{\partial x} & \frac{\partial\phi}{\partial y} & \frac{\partial\phi}{\partial z} \end{vmatrix}$$

$$= \vec{i} \left(\frac{\partial^2\phi}{\partial y \partial z} - \frac{\partial^2\phi}{\partial y \partial z} \right) - \vec{j} \left(\frac{\partial^2\phi}{\partial x \partial z} - \frac{\partial^2\phi}{\partial x \partial z} \right) +$$

$$= 0 \quad \vec{k} \left(\frac{\partial^2\phi}{\partial x \partial y} - \frac{\partial^2\phi}{\partial x \partial y} \right)$$

② Prove that $\text{div}(\text{curl } \vec{F}) = 0$ (or) $\nabla \cdot (\nabla \times \vec{F}) = 0$

Soln:

$$\text{Curl } \vec{F} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ F_1 & F_2 & F_3 \end{vmatrix} \quad \text{if } \vec{F} = F_1\vec{i} + F_2\vec{j} + F_3\vec{k}$$

$$\nabla \times \vec{F} = \vec{i} \left(\frac{\partial F_3}{\partial y} - \frac{\partial F_2}{\partial z} \right) - \vec{j} \left(\frac{\partial F_3}{\partial x} - \frac{\partial F_1}{\partial z} \right) +$$

$$\vec{k} \left(\frac{\partial F_2}{\partial x} - \frac{\partial F_1}{\partial y} \right)$$

$$\nabla \cdot \nabla \times \vec{F} = \left(\vec{i} \frac{\partial}{\partial x} + \vec{j} \frac{\partial}{\partial y} + \vec{k} \frac{\partial}{\partial z} \right) \cdot \left[\vec{i} \left(\frac{\partial F_3}{\partial y} - \frac{\partial F_2}{\partial z} \right) - \vec{j} \left(\frac{\partial F_3}{\partial x} - \frac{\partial F_1}{\partial z} \right) + \vec{k} \left(\frac{\partial F_2}{\partial x} - \frac{\partial F_1}{\partial y} \right) \right]$$

DEPARTMENT OF MATHEMATICS

- ⑦ Find the constants a, b, c so that the vector
 $\vec{F} = (x + 2y + az)\vec{i} + (bx - 3y - z)\vec{j} + (4x + cy + 2z)\vec{k}$
 is irrotational.

Soln:

$$\nabla \times \vec{F} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ x+2y+az & bx-3y-z & 4x+cy+2z \end{vmatrix}$$

$$= \vec{i} \left[\frac{\partial}{\partial y} (4x + cy + 2z) - \frac{\partial}{\partial z} (bx - 3y - z) \right]$$

$$- \vec{j} \left[\frac{\partial}{\partial x} (4x + cy + 2z) - \frac{\partial}{\partial z} (x + 2y + az) \right]$$

$$+ \vec{k} \left[\frac{\partial}{\partial x} (bx - 3y - z) - \frac{\partial}{\partial y} (x + 2y + az) \right]$$

$$= \vec{i} (c+1) - \vec{j} (4-a) + \vec{k} (b-2)$$

Given: \vec{F} is irrotational.

i.e., $\nabla \times \vec{F} = 0$.

$$\vec{i} (c+1) - \vec{j} (4-a) + \vec{k} (b-2) = 0$$

$$c+1 = 0 \Rightarrow c = -1$$

$$4-a = 0 \Rightarrow a = 4$$

$$b-2 = 0 \Rightarrow b = 2$$

- ⑧ Find 'a' so that the vector

$$\vec{A} = (ax^2 - y^2 + x)\vec{i} - (2xy + y)\vec{j}$$
 is irrotational.

Soln:

Given: \vec{A} is irrotational.

$$\nabla \times \vec{A} = 0$$



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Approved by AICTE, New Delhi, Affiliated to Anna University, Chennai

Accredited by NAAC-UGC with 'A++' Grade (Cycle III) & amp;

Accredited by NBA (B.E - CSE, EEE, ECE, Mech & amp; B.Tech.IT)

COIMBATORE-641 035, TAMIL NADU



DEPARTMENT OF MATHEMATICS

$$\nabla \times \vec{A} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ ax^2 - y^2 + x & -(2xy + y) & 0 \end{vmatrix}$$

$$= \vec{i}(0) - \vec{j}(0) + \vec{k}(-2y + 2y) = 0.$$

$\therefore 'a'$ is arbitrary.

- ⑨ Prove $\vec{F} = (y^2 \cos x + z^3) \vec{i} + (2y \sin x - 4) \vec{j} + 3xz^2 \vec{k}$ is irrotational and find its scalar potential ϕ such that $\vec{F} = \nabla \phi$.

Soln:

$$\nabla \times \vec{F} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ y^2 \cos x + z^3 & 2y \sin x - 4 & 3xz^2 \end{vmatrix}$$

$$= \vec{i} \left[\frac{\partial}{\partial y} (3xz^2) - \frac{\partial}{\partial z} (2y \sin x - 4) \right]$$

$$- \vec{j} \left[\frac{\partial}{\partial x} (3xz^2) - \frac{\partial}{\partial z} (y^2 \cos x + z^3) \right]$$

$$+ \vec{k} \left[\frac{\partial}{\partial x} (2y \sin x - 4) - \frac{\partial}{\partial y} (y^2 \cos x + z^3) \right]$$

$$= 0\vec{i} + 0\vec{j} + 0\vec{k}$$

$$= 0$$

$\therefore \vec{F}$ is irrotational.

To find ϕ :

$$\nabla \phi = (y^2 \cos x + z^3) \vec{i} + (2y \sin x - 4) \vec{j} + 3xz^2 \vec{k}$$

$$\text{We know that } \nabla \phi = \vec{i} \frac{\partial \phi}{\partial x} + \vec{j} \frac{\partial \phi}{\partial y} + \vec{k} \frac{\partial \phi}{\partial z}$$

$$\frac{\partial \phi}{\partial x} = y^2 \cos x + z^3 \Rightarrow \phi = y^2 \sin x + z^3 x + f(y, z)$$

$$\frac{\partial \phi}{\partial y} = 2y \sin x - 4 \Rightarrow \phi = y^2 \sin x - 4y + f(x, z)$$

$$\frac{\partial \phi}{\partial z} = 3xz^2 \Rightarrow \phi = y^2 x z^3 + f(x, y)$$

- (10) Show that $\vec{F} = (6xy + z^3) \vec{i} + (3x^2 - z) \vec{j} + (3xz^2 - y) \vec{k}$ is irrotational. Find ϕ such that $\vec{F} = \nabla \phi$.

Soln: $\phi = 3x^2y + xz^3 - yz + c$

- (11) If $\nabla \phi = yz \vec{i} + xz \vec{j} + xy \vec{k}$ then find ϕ .

Soln: $\phi = xyz + c$

- (12) Prove that $\text{div } \hat{r} = 2/r$.

Soln:

$$\text{div } \hat{r} = \nabla \cdot \left(\frac{\vec{r}}{r} \right)$$

$$= \left(\frac{\partial}{\partial x} \vec{i} + \frac{\partial}{\partial y} \vec{j} + \frac{\partial}{\partial z} \vec{k} \right) \cdot \left(\frac{x\vec{i} + y\vec{j} + z\vec{k}}{r} \right)$$

$$= \frac{\partial}{\partial x} \left(\frac{x}{r} \right) + \frac{\partial}{\partial y} \left(\frac{y}{r} \right) + \frac{\partial}{\partial z} \left(\frac{z}{r} \right)$$

$$= \frac{1}{r} - \frac{1}{r^2} \cdot x \frac{\partial r}{\partial x} + \frac{1}{r} - \frac{1}{r^2} y \frac{\partial r}{\partial y} +$$

$$\frac{1}{r} - \frac{1}{r^2} z \frac{\partial r}{\partial z}$$

$$= \frac{3}{r} - \frac{1}{r^2} \left[x \frac{\partial r}{\partial x} + y \frac{\partial r}{\partial y} + z \frac{\partial r}{\partial z} \right]$$

Now $r^2 = x^2 + y^2 + z^2$

$$2r \frac{\partial r}{\partial x} = 2x \Rightarrow \frac{\partial r}{\partial x} = \frac{x}{r}$$