



## DEPARTMENT OF MATHEMATICS

Consider,

$$a_0 x^n \frac{d^n y}{dx^n} + a_1 x^{n-1} \frac{d^{n-1} y}{dx^{n-1}} + \dots + a_{n-1} x \frac{dy}{dx} + a_n y = f(x)$$

→ ①

Where  $a_0, a_1, a_2, \dots, a_n$  are constants and  $f(x)$  is a function of  $x$  is called a homogeneous linear differential equation of order  $n$  with Variable Coefficients.

For this type, substitute

$$z = \log x \quad (\text{or}) \quad x = e^z$$

Then  $x \frac{dy}{dx} = D'y$

$$x^2 \frac{d^2 y}{dx^2} = D'(D'-1)y$$



# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Approved by AICTE, New Delhi, Affiliated to Anna University, Chennai

Accredited by NAAC-UGC with 'A++' Grade (Cycle III) &

Accredited by NBA (B.E - CSE, EEE, ECE, Mech & B.Tech.IT)

COIMBATORE-641 035, TAMIL NADU



## DEPARTMENT OF MATHEMATICS

$$x^3 \frac{d^3 y}{dx^3} = D'(D'-1)(D'-2)y \text{ and so on}$$

Problems:

- ① Transform the equation to constant coefficient equations:

$$x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} + y = \log x \sin(\log x)$$

Soln:

Replace  $\log x = z$

$$x \frac{dy}{dx} = D'y$$

$$x^2 \frac{d^2 y}{dx^2} = D'(D'-1)y$$

Given equation becomes,

$$D'(D'-1)y + D'y + y = z \sin z$$

$$(D'^2 - D')y + D'y + y = z \sin z$$

$$(D'^2 - \cancel{D'} + \cancel{D'} + 1)y = z \sin z$$

$$\boxed{(D'^2 + 1)y = z \sin z}$$