



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Approved by AICTE, New Delhi, Affiliated to Anna University, Chennai

Accredited by NAAC-UGC with 'A++' Grade (Cycle III) &

Accredited by NBA (B.E - CSE, EEE, ECE, Mech & B.Tech.IT)

COIMBATORE-641 035, TAMIL NADU



DEPARTMENT OF MATHEMATICS

Harmonic functions:

An expression of the form $\frac{\partial^2 \phi}{\partial x^2} + \frac{\partial^2 \phi}{\partial y^2} = 0$ is called the Laplace equation in two dimension.

Any function having continuous second order partial derivatives which satisfies the Laplace equation is called harmonic function.

Any two harmonic functions u and v such that $f(z) = u + iv$ is analytic are called conjugate harmonic functions.



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Approved by AICTE, New Delhi, Affiliated to Anna University, Chennai

Accredited by NAAC-UGC with 'A++' Grade (Cycle III) &

Accredited by NBA (B.E - CSE, EEE, ECE, Mech & B.Tech.IT)

COIMBATORE-641 035, TAMIL NADU



DEPARTMENT OF MATHEMATICS

Note:

Both real and imaginary parts of an analytic function are harmonic. But the converse need not be true.

- ① Give an example such that u and v are harmonic but $u+iv$ is not analytic.

Soln:

$$\text{Let } w = \bar{z} = x - iy$$

$$u + iv = x - iy$$

$$\Rightarrow u = x \quad ; \quad v = -y$$

$$\frac{\partial u}{\partial x} = 1, \quad \frac{\partial u}{\partial y} = 0 \quad ; \quad \frac{\partial v}{\partial x} = 0, \quad \frac{\partial v}{\partial y} = -1$$

$$\frac{\partial^2 u}{\partial x^2} = 0, \quad \frac{\partial^2 u}{\partial y^2} = 0 \quad ; \quad \frac{\partial^2 v}{\partial x^2} = 0, \quad \frac{\partial^2 v}{\partial y^2} = 0$$

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 \quad \text{and} \quad \frac{\partial^2 v}{\partial x^2} + \frac{\partial^2 v}{\partial y^2} = 0$$

$\Rightarrow u$ and v are harmonic.

$$\text{But } u_x \neq v_y \quad \text{and} \quad u_y = -v_x$$

$\therefore f(z) = u + iv$ is not analytic.

- ② Prove that $u = e^x \cos y$ is a harmonic function.

Soln:

$$\text{Let } u = e^x \cos y$$

$$\frac{\partial u}{\partial x} = e^x \cos y \quad ; \quad \frac{\partial u}{\partial y} = -e^x \sin y$$

$$\frac{\partial^2 u}{\partial x^2} = e^x \cos y \quad ; \quad \frac{\partial^2 u}{\partial y^2} = -e^x \cos y$$

$$\Rightarrow \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$$

$\therefore u$ is a harmonic function.



DEPARTMENT OF MATHEMATICS

(3) prove that $u = x^2 - y^2$, $v = \frac{-y}{x^2 + y^2}$ are harmonic

but $u + iv$ is not a regular function.

Soln:

$$\text{Let } u = x^2 - y^2$$

$$\frac{\partial u}{\partial x} = 2x \quad ; \quad \frac{\partial u}{\partial y} = -2y$$

$$\frac{\partial^2 u}{\partial x^2} = 2 \quad ; \quad \frac{\partial^2 u}{\partial y^2} = -2$$

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 2 - 2 = 0$$

$\Rightarrow u$ is a harmonic function.

$$\text{Let } v = \frac{-y}{x^2 + y^2}$$

$$\frac{\partial v}{\partial x} = - \frac{[(x^2 + y^2) \cdot 0 - y(2x)]}{(x^2 + y^2)^2} = \frac{2xy}{(x^2 + y^2)^2}$$

$$\frac{\partial^2 v}{\partial x^2} = \frac{(x^2 + y^2)^2 (2y) - 2xy \cdot 2(x^2 + y^2)(2x)}{(x^2 + y^2)^4}$$

$$= \frac{(x^2 + y^2)^2 2y - 8x^2 y (x^2 + y^2)}{(x^2 + y^2)^4}$$

$$= \frac{2y(x^2 + y^2) - 8x^2 y}{(x^2 + y^2)^3}$$

$$= \frac{2y^3 - 6x^2 y}{(x^2 + y^2)^3}$$

$$\frac{\partial v}{\partial y} = - \frac{[(x^2 + y^2) - y \cdot 2y]}{(x^2 + y^2)^2} = \frac{-(x^2 - y^2)}{(x^2 + y^2)^2} = \frac{y^2 - x^2}{(x^2 + y^2)^2}$$

$$\frac{\partial^2 v}{\partial y^2} = \frac{(x^2 + y^2)^2 2y - (y^2 - x^2) 2(x^2 + y^2) 2y}{(x^2 + y^2)^4}$$



DEPARTMENT OF MATHEMATICS

PROBLEMS:

(1) Prove that $w = z^2$ is analytic.

Soln:

We know $z = x + iy$

$$\therefore w = z^2$$

$$= (x + iy)^2$$

$$= x^2 - y^2 + 2ixy$$

$$u + iv = (x^2 - y^2) + i(2xy)$$

$$u = x^2 - y^2 \quad ; \quad v = 2xy$$

$$u_x = 2x \quad ; \quad v_x = 2y$$

$$u_y = -2y \quad ; \quad v_y = 2x$$

$$u_x = v_y \quad \& \quad u_y = -v_x$$

It satisfies CR equations.

$\Rightarrow w = z^2$ is analytic.

(2) Determine whether the function $w = 2xy + i(x^2 - y^2)$ is analytic.

Soln: $w = 2xy + i(x^2 - y^2)$

$$u + iv = 2xy + i(x^2 - y^2)$$

$$u = 2xy \quad , \quad v = x^2 - y^2$$

$$u_x = 2y \quad , \quad v_x = 2x$$

$$u_y = 2x \quad , \quad v_y = -2y$$

$$u_x \neq v_y$$

$$\& \quad u_y \neq -v_x$$

It doesn't satisfy CR equations

$\Rightarrow w = 2xy + i(x^2 - y^2)$ is not analytic.



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Approved by AICTE, New Delhi, Affiliated to Anna University, Chennai

Accredited by NAAC-UGC with 'A++' Grade (Cycle III) &

Accredited by NBA (B.E - CSE, EEE, ECE, Mech & B.Tech.IT)

COIMBATORE-641 035, TAMIL NADU



DEPARTMENT OF MATHEMATICS

$$= \frac{(x^2+y^2)^2 2y - 4y(y^2-x^2)(x^2+y^2)}{(x^2+y^2)^4}$$

$$= \frac{(x^2+y^2) 2y - 4y(y^2-x^2)}{(x^2+y^2)^3}$$

$$= \frac{6x^2y - 2y^3}{(x^2+y^2)^3} = \frac{6}{(x^2+y^2)^3}$$

$$\Rightarrow \frac{\partial^2 v}{\partial x^2} + \frac{\partial^2 v}{\partial y^2} = \frac{2y^3 - 6x^2y}{(x^2+y^2)^3} + \frac{6x^2y - 2y^3}{(x^2+y^2)^3} = 0$$

$\Rightarrow v$ is a harmonic function.

But $\frac{\partial u}{\partial x} \neq \frac{\partial v}{\partial y}$; $\frac{\partial u}{\partial y} \neq -\frac{\partial v}{\partial x}$

$f(z) = u+iv$ is not analytic (or) not regular function.

$u_x = v_y$ $u_y = -v_x$
