



## Stoke's Theorem :

### Statement :

\*If  $\vec{F}$  is any continuous differentiable (vector) function and  $S$  is the surface enclosed by a curve  $C$  then

$$\int_C \vec{F} \cdot d\vec{r} = \iiint_S (\nabla \times \vec{F}) \cdot \hat{n} \, ds.$$

\*Where,  $d\vec{r} = dx\vec{i} + dy\vec{j} + dz\vec{k}$

### Example : 1

Verify Stoke's theorem for a function  $\vec{F} = (y-z+2)\vec{i} + (y-z+4)\vec{j} - 4z\vec{k}$  where  $S$  is the open surface of the cube  $x=0$  to  $x=2$

$y=0$ ,  $y=2$ ,  $z=2$  about the  $xy$  plane



Soln: Given,


$$\vec{F} = (y-z+2)\vec{i} + (yz+4)\vec{j} - 4z\vec{k}$$

Formula:  $\int_C \vec{F} \cdot d\vec{r} = \iint_S (\nabla \times \vec{F}) \cdot \hat{n} \cdot ds$

RHS:

$$\nabla \times \vec{F} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ y-z+2 & yz+4 & -4z \end{vmatrix}$$

$$\begin{aligned} &= \vec{i} \left[ \frac{\partial}{\partial y} (-4z) - \frac{\partial}{\partial z} (yz+4) \right] - \vec{j} \left[ \frac{\partial}{\partial x} (-4z) - \frac{\partial}{\partial z} (y-z+2) \right] \\ &+ \vec{k} \left[ \frac{\partial}{\partial x} (yz+4) - \frac{\partial}{\partial y} (y-z+2) \right] \\ &= \vec{i} (-y) - \vec{j} (1) + \vec{k} (-1) \\ &= -y\vec{i} - \vec{j} + \vec{k} \\ &= 4\vec{i} + \vec{j} + \vec{k} \end{aligned}$$



Surface	Equation	$\hat{n}$	$ds$
$S_1$	$x=0$	$-\vec{i}$	$dydz$
$S_2$	$x=2$	$\vec{i}$	$dydz$
$S_3$	$y=0$	$-\vec{j}$	$dx dz$
$S_4$	$y=2$	$\vec{j}$	$dx dz$
$S_5$	$z=2$	$\vec{k}$	$dx dy$

Along OA  $y=0$ .

$$= - \iint_{OA} (y\vec{i} + \vec{j} + \vec{k}) \cdot (-\vec{i}) dx dz$$

$$= - \int_0^2 \int_0^2 dx dz$$

$$= - \int_0^2 (2) dz$$

$$= - [2(z)^2]_0^2$$

$$= -2(2) = -4$$

Along AB  $x=2$   $\hat{n} = \vec{i}$   $ds = dydz$

$$= \iint_{AB} (y\vec{i} + \vec{j} + \vec{k}) \cdot \vec{i} dy dz$$



$$\begin{aligned}
 &= \int_0^2 \int_0^2 (y) \, dy \, dz \\
 &= \int_0^2 \left( \frac{y^2}{2} \right)_0^2 dz = \int_0^2 \left( \frac{4}{2} \right) dz \\
 &= 2 \int_0^2 dz = 2 (z)_0^2 \\
 &= 2(2) = 4.
 \end{aligned}$$

Along BC:  $y=2$ ,  $\hat{n} = \hat{j}$ ,  $ds = dx \, dz$

$$\begin{aligned}
 &= \int_0^2 \int_0^2 (y\hat{i} + \hat{j} + z\hat{k}) \cdot \hat{j} \, dx \, dz \\
 &= \int_0^2 \int_0^2 dx \, dz = \int_0^2 (x)_0^2 dz \\
 &= 2 \int_0^2 dz = 2(z)_0^2 \\
 &= 2(2) = 4.
 \end{aligned}$$

Along OC:  $x=0$ ,  $\hat{n} = -\hat{i}$ ,  $ds = dy \, dz$

$$\begin{aligned}
 &= - \int_0^2 \int_0^2 (y\hat{i} + \hat{j} + z\hat{k}) \cdot \hat{i} \, dy \, dz \\
 &= - \int_0^2 \int_0^2 y \, dy \, dz \\
 &= - \int_0^2 \left( \frac{y^2}{2} \right)_0^2 dz = - \int_0^2 \frac{4}{2} dz \\
 &= -2 (z)_0^2 = -4.
 \end{aligned}$$

Along CA:  $z=2$ ,  $\hat{n} = \hat{k}$ ,  $ds = dx \, dy$

$$\begin{aligned}
 &= \int_0^2 \int_0^2 (y\hat{i} + \hat{j} + z\hat{k}) \cdot \hat{k} \, dx \, dy \\
 &= \int_0^2 \int_0^2 dx \, dy = \int_0^2 (x)_0^2 dy \\
 &= 2 \int_0^2 dy = 2(y)_0^2 \\
 &= 2(2) = 4.
 \end{aligned}$$



# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



UNIT-I Vector calculus

Stokes theorem

$$\therefore S_1 + S_2 + S_3 + S_4 + S_5 = -4 + 4 + 4 - 4 + 4$$

$$\text{RHS.} = 4.$$

LHS:

$$\vec{F} \cdot d\vec{x} = [(y-z+2)\vec{i} + (yz+4)\vec{j} - 4z\vec{k}] \cdot (dx\vec{i} + dy\vec{j} + dz\vec{k})$$

$$= (y-z+2)dx + (yz+4)dy - 4zdz$$

$$\int_C \vec{F} \cdot d\vec{x} = \int_{S_1} + \int_{S_2} + \int_{S_3} + \int_{S_4} + \int_{S_5} \vec{F} \cdot d\vec{x}$$

On  $S_4$   $x=0$ ,  $dx=0$ ,  $z=0$ ,  $dz=0$ .

$$\int_{S_4} \vec{F} \cdot d\vec{x} = \int_{S_4} (yz+4)dy - 4zdz$$

$$= \int_0^2 (4)dy = 4 \int_0^2 dy$$

$$= 4(y)_0^2 = 8.$$

On OA  $y=0$ ,  $z=0$ ,  $dy=0$ ,  $dz=0$ .

$$\int_{OA} 2dx = \int_0^2 2x = 2(x)_0^2 = 4.$$

On AB  $x=2$ ,  $z=0$ ,  $dx=0$ ,  $dz=0$ .

$$\int_{AB} 4dy = \int_0^2 4dy = 4(y)_0^2 = 8.$$

On BC  $y=2$ ,  $z=0$ ,  $dy=0$ ,  $dz=0$ .

$$\int_{BC} 4dx = \int_0^2 4dx = 4(x)_0^2 = 8.$$

$$\int_C \vec{F} \cdot d\vec{x} = 8 + 8 + 4 + 8 + 8 = 36.$$