



## SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Approved by AICTE, New Delhi, Affiliated to Anna University, Chennai

Accredited by NAAC-UGC with 'A++' Grade (Cycle III) &

Accredited by NBA (B.E - CSE, EEE, ECE, Mech & B.Tech.IT)



Necessary Condition (Cartesian Coordinates) (or)

Cauchy- Riemann equations :-

If the function  $f(z) = u(x,y) + iv(x,y)$  is analytic in a region  $R$  of the  $z$  plane, then

(i)  $\frac{\partial u}{\partial x}$ ,  $\frac{\partial u}{\partial y}$ ,  $\frac{\partial v}{\partial x}$  and  $\frac{\partial v}{\partial y}$  exists

(ii)  $\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$  and  $\frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$  at every point in that region.

Sufficient Conditions:-

If the function  $f(z) = u(x,y) + iv(x,y)$  is analytic in a region  $R$  of the  $z$ -plane if

(i)  $u_x, u_y, v_x$  &  $v_y$  are exists and all are continuous,

(ii)  $u_x = v_y$  and  $u_y = -v_x$ .

Necessary Condition (Polar coordinates) :-

If the function  $w = f(z) = u(r, \theta) + iv(r, \theta)$  is analytic in a region  $R$  of the  $z$ -plane then

i) If  $\frac{\partial u}{\partial r}$ ,  $\frac{\partial u}{\partial \theta}$ ,  $\frac{\partial v}{\partial r}$  and  $\frac{\partial v}{\partial \theta}$  exists.



# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Approved by AICTE, New Delhi, Affiliated to Anna University, Chennai  
Accredited by NAAC-UGC with 'A++' Grade (Cycle III) &  
Accredited by NBA (B.E - CSE, EEE, ECE, Mech & B.Tech.IT)



Problems:-

1) Prove that  $w = z^2$  is analytic.

Sol:- We know that  $z = x + iy$

$$w = z^2 = (x + iy)^2 = x^2 + 2ixy - y^2$$

$$u + iv = (x^2 - y^2) + i(2xy)$$

$$u = x^2 - y^2$$

$$v = 2xy$$

$$u_x = 2x$$

$$v_x = 2y$$

$$u_y = -2y$$

$$v_y = 2x$$

$$u_x = v_y \text{ and } u_y = -v_x$$

It satisfies the Cauchy-Riemann equations.

$w = z^2$  is analytic.

2) Determine whether the function  $w = 2xy + i(x^2 - y^2)$  is analytic.

Sol:-  $w = 2xy + i(x^2 - y^2)$

$$u + iv = 2xy + i(x^2 - y^2)$$



# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Approved by AICTE, New Delhi, Affiliated to Anna University, Chennai

Accredited by NAAC-UGC with 'A++' Grade (Cycle III) &

Accredited by NBA (B.E - CSE, EEE, ECE, Mech & B.Tech.IT)



$$u = 2xy$$

$$v = x^2 - y^2$$

$$u_x = 2y$$

$$v_x = 2x$$

$$u_y = 2x$$

$$v_y = -2y$$

$$u_x \neq v_y \text{ and } u_y \neq -v_x$$

It does not satisfy the Cauchy Riemann equations.

$w = 2xy + i(x^2 - y^2)$  is not analytic.

3) Verify whether  $f(z) = \sinh z$  is analytic using CR equations.

Sol:  $f(z) = \sinh z$

$$\begin{aligned} \sin i\theta &= i \sinh \theta \\ \cos i\theta &= \cosh \theta \end{aligned}$$

$$u + iv = \sinh(x + iy)$$

$$= \frac{1}{i} \sin i(x + iy)$$

$$= \frac{1}{i} \sin(ix + i^2 y)$$

$$= \frac{1}{i} \sin(ix - y)$$

$$= \frac{1}{i} [\sin ix \cos y - \cos ix \sin y]$$

$$= \frac{1}{i} [i \sinh x \cos y - \cosh x \sin y]$$

$$= \sinh x \cos y - \frac{1}{i} \cosh x \sin y$$

$$= \sinh x \cos y + i \cosh x \sin y$$

$$u = \sinh x \cos y \quad u_y = -\sinh x \sin y$$

$$u_x = \cosh x \cos y$$

$$\left[ \frac{1}{i} = -i \right]$$



$$V = \cos h x \sin y$$

$$V_x = \sin h x \sin y$$

$$V_y = \cos h x \cos y$$

$$u_x = V_y \text{ and } u_y = -V_x$$

It satisfies the Cauchy Riemann equations

$f(z) = \sinh z$  is analytic.

4) Show that  $f(z) = |z|^2$  is nowhere analytic.

Sol:  $f(z) = |z|^2$

$$u + iv = x^2 + y^2$$

$$u = x^2 + y^2, \quad v = 0$$

$$u_x = 2x, \quad v_x = 0$$

$$u_y = 2y, \quad v_y = 0$$

$$u_x \neq v_y \text{ and } u_y \neq -v_x$$

It does not satisfy the CR equations.

$\therefore f(z) = |z|^2$  is nowhere analytic.

5) If  $u + iv$  is analytic then  $v - iu$  is also analytic.

Sol:  $u + iv$  is analytic.

i.e., CR equations are satisfied.

$$\text{i.e., } u_x = v_y \text{ and } u_y = -v_x$$

$$\text{i.e., } \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \text{ and } \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$