



Construction of Analytic functions:

Milne-Thomson method

i) To find $f(z)$, when u is given

$$f(z) = \int [\phi_1(z,0) - i\phi_2(z,0)] dz$$

where $\phi_1(z,0) = \left(\frac{\partial u}{\partial x}\right)_{(z,0)}$ and

$$\phi_2(z,0) = \left(\frac{\partial u}{\partial y}\right)_{(z,0)}$$

ii) To find $f(z)$, when v is given

$$f(z) = \int [\phi_1(z,0) + i\phi_2(z,0)] dz$$

where $\phi_1(z,0) = \left(\frac{\partial v}{\partial y}\right)_{(z,0)}$ and

$$\phi_2(z,0) = \left(\frac{\partial v}{\partial x}\right)_{(z,0)}$$

iii) If $u-v$ or $u+v$ is given, then to find

Take $f(z) = u + iv$

if $f(z) = iu - v$

Q. Find the analytic function $f(z)$ whose real part is $u = 3x^2y + 2x^2 - y^3 - 2y^2$

Soln.

Given $u = 3x^2y + 2x^2 - y^3 - 2y^2$

$$\frac{\partial u}{\partial x} = 6xy + 4x$$



$$\phi_1(x, 0) = \left(\frac{\partial u}{\partial x} \right)_{(x, 0)} = 4x$$

$$\frac{\partial u}{\partial y} = 3x^2 - 3y^2 - 4y$$

$$\phi_2(x, 0) = \left(\frac{\partial u}{\partial y} \right)_{(x, 0)} = 3x^2$$

By Milne Thomson method,

$$f(z) = \int [\phi_1(x, 0) - i\phi_2(x, 0)] dz$$

$$= \int [4x - i3x^2] dz$$

$$= \frac{4x^2}{2} - i \frac{3x^3}{3} + C$$

$$f(z) = 2z^2 - iz^3 + C$$

HW: $x^3 - 3xy^2 + 3ix^2y - 3iy^3 + C$
 Ans: $x^2 + 3x^2 + C$

2]. Prove that $v = e^{-x}(x \cos y + y \sin y)$ is harmonic and determine analytic function $f(z)$.

Soln.

Given $v = e^{-x}x \cos y + e^{-x}y \sin y$

$$\frac{\partial v}{\partial x} = [e^{-x} + x(-v)e^{-x}] \cos y + (-1)e^{-x}y \sin y$$

$$= e^{-x} \cos y - x e^{-x} \cos y - e^{-x}y \sin y$$

$$\frac{\partial^2 v}{\partial x^2} = -e^{-x} \cos y - [x(-e^{-x}) + e^{-x}] \cos y + e^{-x}y \sin y$$

$$= -e^{-x} \cos y + x e^{-x} \cos y - e^{-x} \cos y + e^{-x}y \sin y$$

$$\frac{\partial v}{\partial y} = e^{-x}x[-\sin y] + e^{-x}[y \cos y + \sin y]$$

$$= -x e^{-x} \sin y + e^{-x}y \cos y + e^{-x} \sin y$$

$$\frac{\partial^2 v}{\partial y^2} = -x e^{-x} \cos y + e^{-x}[y(-\sin y) + \cos y]$$

$$+ e^{-x} \cos y$$

$$= -x e^{-x} \cos y - y \sin y e^{-x} + e^{-x} \cos y$$

$$+ e^{-x} \cos y$$



$$= -x e^{-x} \cos y - e^{-x} y \sin y + 2e^{-x} \cos y$$

$$\therefore \frac{\partial^2 v}{\partial x^2} + \frac{\partial^2 v}{\partial y^2}$$

$$= -2e^{-x} \cos y + 2e^{-x} \cos y + e^{-x} y \sin y - x e^{-x} \cos y - e^{-x} y \sin y + 2e^{-x} \cos y$$

$$= 0$$

Hence v satisfies Laplace eqn.

$\therefore v$ is harmonic.

By Milne's Thomson method,

$$f(z) = \int [\phi_1(z, 0) + i \phi_2(z, 0)] dz$$

$$\text{where } \phi_1(z, 0) = \left(\frac{\partial v}{\partial y}\right)_{(z, 0)}$$

$$= [-x e^{-x} \sin y + e^{-x} y \cos y + e^{-x} \sin y]_{(z, 0)}$$

$$\phi_1(z, 0) = 0$$

$$\phi_2(z, 0) = \left(\frac{\partial v}{\partial x}\right)_{(z, 0)}$$

$$= [e^{-x} \cos y - x e^{-x} \cos y - e^{-x} y \sin y]_{(z, 0)}$$

$$\phi_2(z, 0) = e^{-z} - x e^{-z}$$

$$\therefore f(z) = \int [0 + i(e^{-z} - x e^{-z})] dz$$

$$= i \left[\int e^{-z} dz - \int x e^{-z} dz \right]$$

$$= i [-e^{-z} - (-x e^{-z} - e^{-z})] + c$$

$$= i [-e^{-z} + x e^{-z} + e^{-z}] + c$$

$$f(z) = i x e^{-z} + c$$

How $v = e^{2x} (y \cos 2y + x \sin 2y)$
 $f(z) = ?$

$$v = x^2 - u^2 + 2i$$



By Milne's Thomson method,

$$F(z) = \int [\phi_1(z,0) - i \phi_2(z,0)] dz$$

$$= \int (e^z + i e^z) dz$$

$$= (1+i) \int e^z dz$$

$$(1+i) F(z) = (1+i) e^z + C$$

$$F(z) = e^z + C$$

5]. If $f(z) = u + iv$ is analytic, find $f(z)$ given that $u + v = \frac{\sin 2x}{\cosh 2y - \cos 2x}$

Soln.

$$\text{Let } f(z) = u + iv \rightarrow (1)$$

$$i f(z) = iu - v \rightarrow (2)$$

$$(1) + (2) \Rightarrow (1+i) f(z) = u + iv + iu - v = (u - v) + i(u + v)$$

$$F(z) = U + iV$$

$$\text{Here } F(z) = (1+i) f(z)$$

$$U = u - v$$

$$V = u + v$$

$$\text{Given } V = u + v = \frac{\sin 2x}{\cosh 2y - \cos 2x}$$

$$\frac{\partial V}{\partial x} = \frac{(\cosh 2y - \cos 2x)(2 \cos 2x) - \sin 2x(-2 \sin 2x)}{(\cosh 2y - \cos 2x)^2}$$

$$\phi_2(z,0) = \left(\frac{\partial V}{\partial x} \right)_{(z,0)} = \frac{(1 - \cos 2x) 2 \cos 2x - 2 \sin^2 2x}{(1 - \cos 2x)^2}$$

$$= \frac{(1 - \cos 2x) 2 \cos 2x - 2(1 - \cos^2 2x)}{(1 - \cos 2x)^2}$$

$$= \frac{(1 - \cos 2x) 2 \cos 2x - 2(1 + \cos 2x)(1 - \cos 2x)}{(1 - \cos 2x)^2}$$



By Milne's Thomson method,

$$f(z) = \int [\phi_1(z, 0) - i\phi_2(z, 0)] dz$$

$$= \int [-\csc^2 z - i(0)] dz$$

$$= -\int \csc^2 z dz$$

$$f(z) = \cot z + C$$

4]. Find the analytic function $f(z) = u + iv$
where $u - v = e^x (\cos y - \sin y)$
Soln.

$$\text{Let } f(z) = u + iv \rightarrow (1)$$

$$if(z) = iu - v \rightarrow (2)$$

$$(1) + (2) \Rightarrow (1+i)f(z) = u + iv + iu - v$$

$$(1+i)f(z) = (u-v) + i(u+v)$$

$$F(z) = U + iV$$

$$\text{Here } F(z) = (1+i)f(z)$$

$$U = u - v$$

$$V = u + v$$

$$\text{Given } U = u - v = e^x (\cos y - \sin y)$$

$$\frac{\partial U}{\partial x} = e^x [\cos y - \sin y]$$

$$\phi_1(z, 0) = \left(\frac{\partial U}{\partial x} \right)_{(x,0)} = e^x [1 - 0] = e^x$$

$$\begin{aligned} \left(\frac{\partial U}{\partial y} \right)_{(x,0)} &= e^x [-\sin y - \cos y] \\ &= -e^x [\sin y + \cos y] \end{aligned}$$

$$\phi_2(z, 0) = \left(\frac{\partial U}{\partial y} \right)_{(x,0)} = -e^x [0 + 1] = -e^x$$



3. Determine the analytic function whose real part is $\frac{\sin 2x}{\cosh 2y - \cos 2x}$

Soln.

Given $u = \frac{\sin 2x}{\cosh 2y - \cos 2x}$

$$\frac{\partial u}{\partial x} = \frac{(\cosh 2y - \cos 2x) \cdot 2 \cos 2x - \sin 2x \cdot (2 \sin 2x)}{(\cosh 2y - \cos 2x)^2}$$

$$\phi_1(x,0) = \left(\frac{\partial u}{\partial x} \right)_{x,0} = \frac{(1 - \cos 2x) \cdot 2 \cos 2x - 2 \sin^2 2x}{(1 - \cos 2x)^2}$$

$$= \frac{(1 - \cos 2x) \cdot 2 \cos 2x - 2(1 - \cos^2 2x)}{(1 - \cos 2x)^2}$$

$$= \frac{(1 - \cos 2x) \cdot 2 \cos 2x - 2(1 + \cos 2x)(1 - \cos 2x)}{(1 - \cos 2x)^2}$$

$$= \frac{2[\cos 2x - (1 + \cos 2x)]}{(1 - \cos 2x)}$$

$$= \frac{-2}{(1 - \cos 2x)}$$

$$= \frac{-1}{\frac{1 - \cos 2x}{2}} = \frac{-1}{\sin^2 x}$$

$$\phi_1(x,0) = -\csc^2 x$$

and $\frac{\partial u}{\partial y} = \sin 2x \frac{\partial}{\partial y} \left[\frac{1}{\cosh 2y - \cos 2x} \right]$

$$= \sin 2x \frac{-1}{[\cosh 2y - \cos 2x]^2} [2 \sinh 2y - 0]$$

$$= \frac{-2 \sin 2x \sinh 2y}{(\cosh 2y - \cos 2x)^2}$$

$$\phi_2(x,0) = \left(\frac{\partial u}{\partial y} \right)_{(x,0)} = \frac{-2 \sin 2x (0)}{(1 - \cos 2x)^2} = 0$$



$$\begin{aligned}
 &= \frac{2 \cos 2x - 2(1 + \cos 2x)}{1 - \cos 2x} \\
 &= \frac{2[\cos 2x - 1 - \cos 2x]}{1 - \cos 2x} \\
 &= \frac{-2}{1 - \cos 2x} = \frac{-1}{\frac{1 - \cos 2x}{2}} = \frac{-1}{\sin^2 x}
 \end{aligned}$$

$$\phi_2(x, 0) = -\csc^2 x$$

$$\begin{aligned}
 \frac{\partial v}{\partial y} &= \frac{\sin 2x - 1}{(\cosh 2y - \cos 2x)^2} [2 \sinh 2y - 0] \\
 &= \frac{-2 \sin 2x \sinh 2y}{(\cosh 2y - \cos 2x)^2}
 \end{aligned}$$

$$\phi_1(x, 0) = \left(\frac{\partial v}{\partial y} \right)_{(x, 0)} = 0$$

By Milne's Thomson method,

$$\begin{aligned}
 F(z) &= \int [\phi_1(z, 0) + i \phi_2(z, 0)] dz \\
 &= \int [0 + i(-\csc^2 x)] dz \\
 &= -i \int \csc^2 x dx
 \end{aligned}$$

$$F(z) = i \cot z + C$$

$$(1+i)f(z) = i \cot z + C$$

$$f(z) = \frac{i}{1+i} \cot z + \frac{C}{1+i}$$

$$= \frac{i(1-i)}{(1+i)(1-i)} \cot z + C$$

$$= \frac{i - i^2}{1 - i^2} \cot z + C$$

$$f(z) = \frac{(1+i)}{2} \cot z + C$$